

Papain-Volume 1

1

525



FOOD AND DRUG
Research LABORATORIES, INC.

Vol. I

525

R E P O R T

ACCESSION NO. 462

PAPAIN

CAS. REG. NO. PM9001734

Submitted to:


GRAS Review Branch (BF-335)
Bureau of Foods
Food and Drug Administration
200 C Street, S.W.
Washington D.C., 20204

Attention:

Mr. Alan Spiher,
Project Manager

Date: January 28, 1974

Laboratory No: 1216


James Bond, Manager
Monograph Group



Howard Feinman, Director
Biological Sciences

TABLE OF CONTENTS

| <u>SUBJECT</u> | |
|------------------------------|-----|
| Summary | 1 |
| Chemical Information | 5 |
| Biological Data | |
| Acute and Short Term Studies | 9 |
| Special Studies | 18 |
| Bibliography | B-1 |

Summary

Papain is obtained from the fruit and leaves of the carica papaya plant and contains enzymes similar to pepsin, but acts in both neutral and alkaline medium. The potency of papain varies according to the method of preparation, and the usual grade digests about 35 times its weight of lean meat. The best grades are claimed to render soluble 200 to 300 times their weight of coagulated egg albumin in an alkaline medium. A temperature range from 60 to 90°C is favorable for the digestive process, and 65°C is regarded as the optimum temperature, with the pH at 5.0. Medically, papain has been used to prevent adhesions and sloughing in infected wounds, but allergic manifestations may occur (351a).

Meat tenderizing by the ultimate consumer is usually accomplished by sprinkling a meat cut with a powdered enzyme preparation, and a typical composition for this surface treatment contains 2% commercial papain. The most recent development in the enzymatic meat tenderization is the injection of proteolytic enzyme solutions into the vascular systems of cattle before slaughter. The vascular system effectively distributes the proteolytic enzyme throughout the tissues. Ante-mortem tenderization is now practiced on a fairly large scale by meat packers using low concentrations of specifically purified papain which is equivalent to 5 to 30 parts/million of commercial papain, based on the total weight of the animals. To avoid overtenderization, the meat must be refrigerated until used, and this ante-mortem tenderization

permits the production and sale of a much higher percentage of tender cuts from all grades of beef and particularly from the lower grades (172a).

When injected intracerebrally in mice, .03 ml of a solution containing 0.0005 g of papain/10 ml produced chronic spasms which invariably progressed to death within 7 to 10 minutes; given orally to rats, 200 to 600 mg of papain produced an analgesic and anti-inflammatory activity which was similar to acetylsalicylic acid (521, 134).

A 2% Vermizym in Ringer's solution perforated the skin of round worms within 2 to 5 hours and digested the body of the worms within 4 to 5 hours. A dose of 5 tablets, administered 5 times, at hourly intervals was effective in removing round worms from the human body but was not sufficient to totally erradicate the eggs. Oral administration of Vermizym produced little toxic effects, and the LD₅₀ in mice was 12.5 to 18.5 g/kg and orally in rats, more than 10 g/kg. Intravenous injection into guinea pigs of 50 mg/kg of Vermizym killed the animals due to respiratory arrest. Intravenously, 60 mg/kg in rats caused a decrease in the erythrocyte and leukocyte counts, but oral administration to 10 g/kg did not have any effect on the erythrocyte nor leukocyte counts. The administration caused erosion and hemorrhage to the stomach and intestinal mucosa, but contact with the powder preparation or a concentrated solution of Vermizym for 1 hour did not produce marked damage to the dog stomach or intestinal

mucosa. Vermizym had a more damaging effect on the conjunctiva and the abraided skin than on the stomach and intestinal mucus membrane. The actions of Vermizym described above disappeared when the preparation is heated at 80°C for more than 5 minutes (508).

In several case reports it was found that the use of papain solutions on impactions in the esophagus resulted in death, and when studied under experimental conditions in dogs, it was revealed that there was a destruction of esophageal tissues. These results led to the public "urge of extreme caution in the use of papain solutions for treatment of meat impaction in the esophagus" (11a, 225).

Due to the digestive properties of papain, calcium oxalate and calcium ammonium phosphate stones were used to study the enzymatic dissolution of kidney stones. It was found that a solution of 100 mg of papain produced a 24 to 26% weight loss of the oxalate stones and a 56 to 62% weight loss of phosphate stones after 96 hours of incubation (276).

Inhalation of intratrachea administration of papain μ g to rats, dogs and hamsters produced an emphysema which resembled that observed in humans. In the rat the lesions showed neither regression or resolution beyond the first 30 days following a 4-day exposure (288, 268, 407).

When papain was administered through the rabbit ear vein, it was observed that the permeability of the articular cartilage decreased and that the hexosamine and uronic acid content of the ear

cartilage was significantly decreased (305, 187, 59, 79).

Papain-containing solutions which came in contact with the rat salivary glands produced hypertrophy of the glands, but when administered directly into the stomach, hypertrophy was not observed indicating that the papain must come in direct contact with the gland. It was also demonstrated that papain significantly stimulated acid formation in saliva (565, 143, 161).

The gut content of the animals fed diets containing papain or fed diets in which papain was administered soon thereafter showed that papain increased the digestion of protein in the stomach (583, 18, 221).

PAPAIN

Chemical Information

I. Nomenclature

A. Common Names (351a)

Vegetable pepsin

Papase

B. Chemical Names

No chemical names were encountered in the literature searched.

C. Trade Names (11a, 351a)

Nematolyt

Airbuz

Vermizym

Summetrin

Tromasin

Caroid

Velardon

Papayotin

SL Papain type A-300, purified papain.

D. Chemical Abstracts Unique Registry Number

PM 9001-73-4

II. Empirical formula

No empirical formula was encountered in the literature searched

III. Structural Formula

No structural formula was encountered in the literature searched.

IV. Molecular Weight (533a)

20,000

27,000 \pm 2,000

20,289

V. Specifications

A. Chemical

No chemical specifications were encountered in the literature searched.

B. Food grade (158a)

Assay. Not less than 6000 N.F. Units of papain activity per mg. (Note: One N.F. Unit of papain activity is the activity which releases the equivalent of 1 mcg. of tyrosine from a specified casein substrate under the conditions of the Assay, using the enzyme concentration which liberates 40 μ g. of tyrosine per ml. of Standard Solution.)

Loss on drying.

Not more than 7%.

pH of a 1 in 50 solution

Between 4.8 and 6.2

Limits of impurities:

Arsenic (as As).

Not more than 3 parts per million (0.0003%)

Heavy metals (as Pb).

Not more than 10 parts per million (0.001%).

C. Official Compendia

Food Chemical Codex

VI. Description (158a)

A. General Characteristics

It occurs as a white to light tan amorphous powder or as light brownish gray to weak reddish brown granules with a characteristic odor and taste.

B. Physical Properties

It is partially soluble in water, the solution being colorless to light yellow and somewhat opalescent. It is practically insoluble in alcohol, in chloroform, in ether and most organic solvents; slightly hygroscopic; partially soluble in glycerol.

C. Stability in Containers

Store in tight, light-resistant containers in a cool, dry place. Significant loss of activity resulted when heated to 60° for 30 minutes. Very stable at 30° between pH 5 and 7.

VII. Analytical Methods (43)

Papain

Title:

The determination of the concentration of hydrolytic enzyme solutions: -Chymotrypsin, trypsin, papain, elastase, subtilisin and acetylcholinesterase.

Principle Determination Step:

Titration

Principle Separation Techniques:

None

Principle Analytical Reagent:

p-Nitrophenyl N-Benzoyloxycarbonyl-L-tyrosinate.

Moiety Measured:

Active cites

Substrate:

**p-Nitrophenyl N-benzyloxcarbonyl-
l-tyrosinate.**

VIII. Occurrences and Levels

A. Plants

Bark of trunk, immature fruit, and leaves of *Carica papaya* Linne (Fam. Caricacea) 1 pound dried latex per tree per year.

B. Animals

No occurrences or levels in animals were encountered in the literature searched.

C. Synthetics

No occurrences or levels in synthetics were encountered in the literature searched.

D. Natural Inorganic Substances

No occurrences or levels in natural inorganic substances were encountered in the literature searched.

To study toxicity of intracerebrally injected papain in mice, .03 ml of a solution containing .00020, 0.00010 or 0.00005 g of papain/10 ml were administered. Within 3 to 5 minutes, there were generalized chronic spasms which invariably progressed to tonic spasms and death within 7 to 10 minutes after injection (521).

Studies using Manor Farms female rats weighing 90-140 g showed that 200 to 600 mg doses of papain have oral analgesic and anti-inflammatory activity which was not significantly different from that of acetylsalicylic acid (134).

"Vermizym", a papain preparation with enhanced anthelmintic effectiveness and stability, was studied to determine some of the pharmacologic and toxicological properties. To study the in vitro digestion of pig round worms, one round worm each was placed in 30 cc of a 2% Vermizym-Ringer emulsion and was kept at 37° C. The results show that in 1 1/2 hours white erosion spots were observed on the surface of the worms, and that after 2 1/2 hours digestion of the worms proceeded rapidly. In 5 hours, the external body of the worm disappeared, but the worm was still capable of spontaneous movement. When the same procedures as described were performed using solutions of Vermizym which had been heated to 65°, 70°, 75°, or 80° for 5 minutes and cooled immediately to 37°, the emulsions preheated to 80° or above for 5 minutes completely lost their anthelmintic capacity. It was also found that there was a decrease in the anthelmintic effectiveness, dependent upon the preheating temperature. The changing of the

pH of a Vermizym-Ringer emulsion showed that the anthelmintic activity was completely lost at pH below 3 or above 9. When compared with papayotin, it was found that Vermizym produced perforation in the worm body in 2 to 3 hours, while within 5 hours papayotin only produced erosion of the skin (508).

Five men patients aged 27 to 58 years old, who were infested with round worms, were administered before breakfast 5 tablets of Vermizym at hourly intervals for 5 consecutive hours; round worms and round worm eggs found in the stools were collected during the following 7 days. Four of the 5 patients passed half-digested round worms in their stools, while no side affects were noted in any of the patients nor were there any abnormalities of the stools excreted (508).

Fasted male mice weighing 10 to 15 g were administered orally 0.25 to 0.75 cc/10 g of body weight of a 25 or 50% Vermizym emulsion. The LD₅₀ was 12.5 to 18.7 g/kg and the symptoms observed were pilo-erection, difficulty in breathing, diarrhea, and weakness. Death was due to respiratory arrest. There was no difference in the toxicity due to the different concentrations, and animals which survived 1 to 2 days had blood in the stool, swelling, erosion, and hemorrhage of the mucus membranes of the anus. Gross autopsy revealed small hemorrhage spots of the serous membrane of the small intestines especially the jejunum, and in some it was observed that the intestinal walls had become thinner. No other distinct gross changes were observed in other organs (508).

1 to 5% solutions of Vermizym in 0.9% saline was injected into the upper leg vein of the guinea pig at a rate of 0.5 cc/minute. Two of 4 animals died after the administration of 50 mg/kg, and 4 of 5 died after the administration of 100 mg/kg and death was due to respiratory arrest. The animals were opened immediately after death, and the lungs showed marked expansion while the heart continued to beat. When the solution was preheated to 80° C for 5 minutes and injected intravenously into the guinea pig, a dose as high as 300 mg/kg failed to produce any toxic symptoms (508).

The same solution as described in the above paragraph was injected into the rat tail vein at a dose of 60 mg of Vermizym/kg. The results showed that there was a marked decrease in the number of red blood cells and white blood cells, and it was also shown that when the solution was administered orally to the rats at an equivalent of 10 g of Vermizym/kg, the rat was not killed, and the blood cell count did not decrease (508).

A rabbit eye irritation study was conducted in which 0.2 cc of a 10% or 25% Vermizym emulsion in 0.9% saline or 0.05 g of powder was applied to the conjunctiva of rabbits. One minute after the application of the appropriate solution, the eyes were washed with physiological saline, and the conjunctiva and cornea were examined at varying time intervals. The results showed the eyes to be closed, tearing, and bloodshot, with discharges; swelling or turbidity in the cornea could not be seen (508).

To the abraided and unbraided skin of rabbits 25% or 50% Vermizym emulsion in 0.9% saline were applied for a period of 1 hour. In the unbraided skin mild hemorrhagic spots were observed when treated with a 50% solution and in the skins of the braided animals 25% or 50% solutions showed blood in the solutions within several minutes after the treatment. At the end of 1 hour in the abraided animals there was a strong congestion and hemorrhage at the site of application. When the Vermizym emulsion was preheated at 80°C the symptoms were not observed (508).

When 0.3 cc of 5%, 10%, or 25% Vermizym emulsion in 0.9% saline was injected subcutaneously into the shaved backs of guinea pigs, it was found that with concentrations as low as 5% the site of injection gradually turned dark purple and became necrotic. The same changes were observed at the higher concentration but were found to be more rapid and the area involved much larger (508).

0.3 cc of a 25% Vermizym emulsion and 0.9% saline was injected into the thigh muscle of dogs and shortly after the injection the dogs refused to walk and lifted their legs expressing pain but within 3 hours they started to walk normally. With a 50% emulsion the symptoms were as above but there was also marked local swelling and the symptoms of pain did not disappear until 18 hours after the injection (508).

A 6 cm portion of the small intestines of dogs was used to study the action of Vermizym on the intestinal mucus membrane. 3 cc of a 50% Vermizym emulsion was injected into the small intestines and in some cases 1.5 g of Vermizym powder was placed inside the intestines; it

was found that in 1 hour after the application of either emulsion or powder the mucus membrane was congested and swollen and after 3 hours some dogs had detachment and erosion of the epithelium (508).

Fasted dogs were administered 5 cc of 20% Vermizym emulsion/kg to study the action on the stomach mucosa. Five hours after the medication the animals were sacrificed and the stomach mucosa examined. Gross examination did not reveal any damages. Further study of the action on the stomach mucosa showed the same dose as above of Vermizym administered at 5 and 24 hours to dogs preheated with hexylresorcinol, was without affect on hexylresorcinol induced impairment (508).

5% Vermizym in 0.9% saline was injected into the leg vein of the dog at a dose which was equivalent to 40 to 50 mg/kg to study the effects on blood pressure. The results showed a marked decrease in the arterial blood pressure and a slight increase in the portal vein pressure and repeated injections into the leg vein produced tachyphylaxis (508).

Application of more than 5×10^5 of Vermizym to the guinea pig intestinal tract induced slower contractions. Repeated treatments resulted in a gradual decrease in the degree of the drug affect on the contraction of the intestines. These affects of Vermizym continued to be observed in the presence of atrophine sulfate and neoantergan maleate. The responses as described for the guinea pig intestinal tract were also observed in the virgin rat uteri (508).

Rabbits and dogs were injected intravenously with 60 to 100 mg papain in 5 ml of vehicle and 350 to 400 mg papain in 20 ml of vehicle, respectively, to investigate the potential coagulation defect. Studies demonstrated that there was an immediate release of heparin after injection of papain, and the blood of both species became incoagulable. In rabbits the incoagulability persisted for many hours, and, in some cases, a mild disturbance of the coagulation factor was still observed 2 weeks after the injection (358).

To determine the local and general action of basic proteins, two studies were performed on rabbits, one in which 30 mg/kg of papain was administered intravenously, and one in which 50, 100, and 200 μ g were injected subcutaneously. Papain, administered intravenously, produced a progressive increase in the number of leukocytes, which peaked 2 and 3 hours after injection, and returned to normal within 6 hours. When 100 mg papain was administered subcutaneously, edema, hyperemia and leukocytosis were observed within 2 1/2 hours after injection (89).

Rabbits were used to determine the role of papain in hemato-
poresis. Three dosage schedules were followed with doses of 50, 75, 100, and 150 mg/day, orally administered. The second group of rabbits received daily oral doses of a mixture consisting of 200 mg papain and 50 mg of ascorbic acid. After a 27-day period of medication with papain alone, the animals exhibited a stable condition of hypochromic anemia. After the 27th day papain was added to the normal diets of these rabbits, and there was regeneration of blood pigment

in all animals. It was observed that the rate of hemoglobin regeneration did not appear to be proportional to the dose of papain administered, as the response was observed in practically all dose levels. In the group in which papain and ascorbic acid was administered, it was found that at the end of 9 days of treatment the rate of hemoglobin regeneration was much higher than control. After 10 days of treatment, the hemoglobin level was increased well above the controls in groups which had received a papain, ascorbic acid, and iron mixture. The major finding in these 3 studies was that papain has a certain hematopoietic activity and that this activity is significantly greater when papain is given in association with some other factors present in the mixture (62).

Calcium oxalate and calcium ammonium stones were models used to study the enzymatic dissolution of kidney stones. The stones were placed in a solution of 10 ml urine + 100 mg of papain and the weights of the stones were determined on an analytical scale after 24, 48, 72, and 96 hours. The results show that treatment with urine-papain solution caused a 24 to 26% weight loss of the oxalate stones and a 56 to 62% weight loss of the phosphate stones after 96 hours of incubation (276).

Pigeons and isolated frog hearts were used to study the pathological changes in tissues and organs. In some pigeons, the administration of papain hastened emesis. When the isolated frog hearts were irrigated

with 1 to 2% papain, it strengthened the contractions of both auricle and ventricle by lengthening the refractory period, diminishing fibrillation and other cardiac irregularities, but it had no effect upon the amplitude of the heartbeat (5).

A case report and experimental study on dogs was reported with regard to perforation of the esophagus after the use of Caroid, a proprietary decongestant containing papain. A 27 year-old female was admitted to the hospital with an esophageal perforation. Ten days earlier, a bolus of meat had lodged in the cervical section of her esophagus. A solution of caroid had been administered to digest the meat. Twelve hours after the administration of the Caroid, the patient was found in a state of shock, exhibiting subcutaneous emphysema in the neck. Two days after hospitalization massive hemorrhage suddenly developed from the cervical mediastinotomy, and the patient died. Autopsy revealed that the walls of the common carotid artery were eroded and that approximately 2 inches of the cervical portion of the esophagus were completely destroyed. These findings suggested a study to determine the role of caroid in the digestion and perforation of the patient's esophagus. A bolus of meat was impacted into the esophagus of anesthetized dogs, and a caroid solution was introduced through a polyethylene catheter. Experiments with control animals demonstrated that meat alone produced no damage, but the caroid-treated animals exhibited some degree of damage to the esophageal wall, indicated by reddening of the mucosa and thickening of the wall at the side of the enlarged bolus. In addition, the lungs of each

caroid-treated dog exhibited edema which, in some instances, was sufficient to cause rapid death. "The results of these findings urge extreme caution in the use of Caroid for treatment of meat impaction in the esophagus" (11a).

After papain was used to soften and dislodge impacted meat from an esophagus, a patient suffered extensive destruction of the esophageal wall. Her physicians believed that when papain contacts a neucrotic area in the esophageal wall, it may induce further neucrosis in the adjacent tissue which, in turn, leads to extensive destruction of the esophagus and surrounding structures; therefore, they felt the use of papain is contraindicated in the treatment of impacted meat in the esophagus (225).

Fasting rat studies were done to determine the effect of the activity state of papain in in vivo activity. Milk clotting, egg digestion, and gelatin digestion were used to establish the in vivo activity of several papain preparations; these preparations were administered, along with a test meal, to rats in doses ranging from 1.25 to 20 mg. Data obtained in these studies revealed that papain increased the digestion 1.5- to 2.1-fold and that the gastric contents did not adversely effect the digestive activity of papain (221).

To determine whether papain increases the digestion of protein in vivo, young adult rats were fed a diet containing 0.5 g papain/100 g total diet for 7 to 15 minutes. The animals were sacrificed 2.5 hours after being fed the diet, and their stomach contents removed. An

analysis of these contents revealed that there was a 1.5 to 2.1 greater amount of protein digestion produced by feeding papain (18).

Mongrel dogs were fed soybean diets supplemented with 0.2% papain to determine the in vivo proteolytic activity of papain as demonstrated by nitrogen balance. The results revealed that the addition of papain to the diet significantly increased the rate of digestion of soybean protein, and that the papain addition also significantly reduced the amount of soybean protein required to maintain the dogs' nitrogen equilibrium (583).

Healthy male subjects ranging in age from 21 to 25 years who were given 7.5 mg of papain orally every half-hour to a total of 6 doses, showed a reduction of fibrin formation time but prolonged clot formation (255).

Male Sprague-Dawley rats, weighing 150 to 200 g, were exposed to an aerosolized 10% solution of papain in isotonic saline to study papain-induced emphysema during a 4-hour exposure interval. At least 2 animals were sacrificed at 0, 1, 2, 3, 4, 6, 8, 12, 19, and 24 hours and at 2, 3, 4, 14, 33, 16, and 90 days after exposure. The results demonstrated that the inhalation of papain produced pulmonary lesions which resembled centrilobular emphysema, and these lesions showed neither regression nor resolution beyond the first 30 days following the 4-day exposure (268).

Morphological and biochemical assessment of papain-induced emphysema were conducted on 6-to 8-week-old male and female Syrian hamsters,

weighing between 90 and 120 g. 1 mg of papain/100 g of body weight was administered intratracheally to the hamsters, and animals from control and medicated groups were sacrificed at 1/2, 1, 2, 4, 6, 8, 32, hours and at 4 days. The results show that alveolar departitioning occurred quickly after a single dose of papain, and after 8 days, capillary obliteration was observed. The authors felt it important to note that a single dose of proteolytic enzyme causes collagen to disappear from the alveolar wall (288).

In a study on experimental papain-induced emphysema, mongrel dogs weighing between 24 and 69 pounds were intratracheally administered 1 or 4 injections of 1 or 2 mg papain/pound during a period ranging from 7 to 28 days. The data obtained from these studies showed that there were no significant changes in the arterial blood gas tension, acid base balance, or pulmonary resistance. However, the defusing capacity decreased and the total lung capacity, functional residual capacity, and residual volume increased, while elastic recoil decreased. Necropsy revealed that the resistance of airways smaller than 2 mm in diameter was increased, while resistance of larger airways decreased, and the condition induced by papain was similar both pathophysiologically and morphologically to human panlobular emphysema (407).

Nine-week-old albino rabbits weighing 1 kg were used to study the immunosuppressive activity of the 3a-1 fraction of papain. The results showed that when 10 mg of papain were administered intravenously (followed within 12 to 18 hours by antigen inoculations), there was a

marked immunosuppression. However, when papain was given 8 to 10 days after the antigen inoculation, normal antibody titers occurred. It was also observed that when papain inoculations preceded the antigen injections by 10 to 12 days, there was a reduction in antibody titers (101).

The effect of age and papain on certain steroid hormones was studied in reference to the permeability of articular cartilage in adolescent and adult New Zealand rabbits. A 1.5% solution of papain was injected intravenously by a constant infusion pump into the marginal ear vein at a rate which would deliver 35 mg/kg over a 15 minute interval. Comparison of results between adolescent and adult rabbits revealed that the permeability of the articular cartilage decreased the permeability index by 10.25, a 25% difference from the untreated group. Adult rabbits did not respond to the injections. Further, in the adult animals, papain had no effect on the permeability of cartilage in those animals pretreated with cortisone (305).

To determine the chemical changes in rabbit aorta and ear cartilage induced by papain and calciferol, rabbits were given a 1% solution of papain intravenously on the 1st, 4th, 7th, and 10th day, or papain in conjunction with calciferol given orally for 10 consecutive days. The results showed that papain significantly lowered the hexosamine and uronic acid contents of ear cartilage, that in the papain-treated animals, there was a decrease in aortic hexosamine and uronic acid, and that when given simultaneously with calciferol, there was no effect on the extent of mineralization (187).

Male white rabbits, weighing 2 to 3 kg, were injected in the ear vein with 30 mg papain/kg to determine the effect of intravenous papain on cartilage glycosaminoglycans. One animal each was sacrificed at 0, 2, 6, 17.5, 18, 24, 48, and 72 hours and its ears removed immediately. The results showed that there was a decrease of non-sulfated and sulfated glycosaminoglycans in rabbit ear cartilage following the intravenous injection of papain with the maximum effect occurring at 24 hours (59).

New Zealand albino rabbits, weighing approximately 1 kg, were injected intravenously in the ear vein with .75 ml of a crude papain solution to determine the release of chondroitin sulfate from rabbit cartilage. Blood and urine samples were analyzed, and revealed that following the injections, blood and urine contained a mucopolysaccharide resembling chondroitin sulfate (in chemical and physical properties). Although the gross and histological changes were not described, it was stated that the time of appearance of the mucopolysaccharides closely correlated with changes in the ear and other organs of the animals (79).

A study of the relationship of serum cholesterol level, aorta cholesterol content and the degree of atherosclerosis in experimentally induced cholesterol-atherosclerosis was conducted in rabbits. The addition of papain to cis-oleic acid and cholesterol containing diets increased the serum cholesterol levels by half, and in animals fed cis-oleic acid alone the amount of cholesterol content in the aorta was increased 2-fold without evidence of severe aortic lesions (349).

Male Holtzman rats, 60 to 90 days of age, were used for 3-to 4-day studies of the taste receptors sialadenotrophic action of proteolytic enzymes. In the first series of tests, the proteolytic enzymes were administered as dietary additives and in the second series, 400 mg/day of papain were administered by medicine dropper and stomach tubing. At the end of the experiments, the submaxillary glands were removed. These studies demonstrated that the papain preparation administered by medicine dropper showed a significant increase ($p < .01$) in the weight of the glands but was not effective when administered by stomach tube. This difference suggested that papain must come into contact with the oral cavity in order to produce an effect on gland weight (565).

In another series of studies, sympathectomy (extirpation of the superior cervical ganglion) and parasympathectomy (section of the chorda tympani nerve) were performed by surgical procedures, and animals were fed for 5 days a diet which contained 1% papain by weight. Pre-ganglionic parasympathectomy resulted in a significant ($p < .01$) reduction in the weight of the ipsilateral gland, while post-ganglionic sympathectomy had only a slight effect (if any) on the ipsilateral glandular weight in control animals. The addition of papain to the diet resulted in a marked increase in the weight of the glands. In the animals having unilateral sympathectomy, the response of the ipsilateral gland to papain was markedly reduced. On the other hand, the parasympathectomized gland still maintained most of its response to papain; after

complete denervation, the response was absent. This result indicated that the maximum sialadenotrophic action can only occur if both autonomic branches are intact (565).

In another study, one week after resection of glossopharyngeal nerves or lingual nerves, 200 mg of papain were given twice daily to one group of rats, while in the second group, papain administration began 24 hours after the surgery. The duration of papain administration in both experiments was 3 days, and the data obtained showed that the usual increase in the weight of the glands followed the intraoral administration of papain was completely absent when the nerves were sectioned (565).

Male Long Evans and Holtzman rats, weighing between 50 and 62 g, were used to study submaxillary gland hypertrophy after diets with proteolytic enzymes. The animals were sacrificed after 12 days on a diet containing 10% papain, and it was found that the weight of the submaxillary gland was significantly decreased, and the weight of the salivary gland was increased. Histologically, the submaxillary gland showed marked enlargement of the alveoli with pale mucoid-appearing hypertrophied acinar cells, abundant PAS-staining cytoplasmic granules, and a peripherally displaced, disfigured, and hyperchromatic nucleus (143).

Saliva was utilized to study the effect of proteolytic enzymes on acid formation in mice, and it was found that papain significantly stimulated acid formation (161).

In vivo and in vitro studies were conducted on rats and/or rat tissues to determine the level and mode of activation of peptidase in liver, spleen, and kidney homogenates, and their relationship to orally-given proteases. In the in vitro studies, animals were given 200 mg of papain by stomach tube every half-hour for a period of 1 1/2 hours and were then sacrificed. Within 30 minutes after oral administration of papain, the trypsin-like activity, leucine, aminopeptidase activity and carboxypeptidase activity in the liver, spleen, and kidney were significantly elevated (254).

In a letter to the editor, it was stated that when human lymphocytes were incubated for 144 hours in a medium containing 8 to 40 µg/ml of papain, there was a higher incidence of blastic elements than found in the controls demonstrating a mitogenic action of papain (344).

The mitotic changes in liver after intraperitoneal papain protease injection were studied in male Wistar rats weighing 110 to 130 g. The animals were given single injections of enzyme solutions showing units of caseinolytic activity of 0.05, 0.1, 0.25, 0.4, and 0.6 in a volume of 0.25 ml/100 g of body weight and were sacrificed 48 hours after injection. There was a marked increase in mitotic index but no pathological alterations in the liver tissue, and observations of the kidney, heart, small intestines, and adrenal also revealed no mitotic changes (284).

During an 18-to 21-week study, it was demonstrated that the fatty liver observed in insulin-treated depancreanized dogs could be pre-

vented by the daily administration of the plant proteolytic enzyme papain (153).

In sheep, papain was found to be a successful anthelmintic against *hamonchus contortus*. It demonstrated some advantages over other anthelmintics as it was not toxic to the host and was administered safely to young, sick, or weak animals and to animals in advance stages of pregnancy (127).

PAPAIN

Bibliography

** Article to be found in summary

* Article to be found in text

PAPAIN

Bibliography

- (1) Abbott, D. C., et al. 1952. A spectrophotometric modification of the Ayre-Anderson method for estimation of proteolytic activity. Arch. Biochem. and Biophys. 38: 85-90
- (2) Abelin, I., and H. Pfister. 1953. Determination and activation of papain. Z. Physiol. Chem. 295: 323-332
- (3) Abeyaratne, M. R., et al. 1964. Effect of glycerol on the resistance of blood and plasma clots to the action of proteolytic enzymes. Nature 202(4927): 98-99
- (4) Agren, G. 1944. Peptidase activity in papain preparations. Acta Physiol. Scandinav. 7: 354-361
- * (5) Ahmann, C. F. 1935. Pathologic changes in tissues and organs affected by deficiency diseases or by toxic substances. Fla. Agr. Expt. Sta., Ann. Rept. 72(1936)
- (6) Aho, K., and C. L. Christian. 1966. Incomplete antibodies. I. Effect of papain on red cells. Blood 27(5): 662-669
- (7) Alessandrini, M. E. 1947. Activity of papain and of papayotin. Rend. ist. super. sanita. 10: 517-522
- (8) Ambros, O., and A. Harteneck. 1928. Über die Wirkungen von Proteasen pflanzlicher Milchsäfte. Unters. über Enzyme 2: 1698-1707
- (9) Ambrus, J. I., et al. 1957. Clinical and experimental studies on fibrinolytic enzymes. In: Proteolytic enzymes and their clinical application. Ann. New York Acad. Sci. 68(1): 97-137
- (10) Annon, R., et al. 1953. Activated papain and eggs of nematodes. Med. Monatsschr. 7: 705-708

- (11) Anderson, A. J., et al. 1963. Glycoprotein excretion following the injection of papain into rabbits. *Nature* 197: 707
- * (11a) Anderson, H. A., et al. 1959. Perforation of the esophagus after the use of a digestant agent: report of case and experimental study. *Ann. Otol.* 68: 890-96
- (12) Anghileri, L. J. 1968. Colloidal chromic phosphate; an improved method of preparation and in situ precipitation by injection of papain. *Int. J. Appl. Radiat. Isotopes* 19(4): 415-420
- (13) Annetts, M. 1936. The digestion products formed by the action of papain on egg albumin. *Biochem. J.* 30: 1807-1814
- (14) Anonymous. 1970. Some observations on the effects of crude papain on tenderization of poultry meat. *J. of Food Science and Technology* 7(1): 40-41
- (15) Anson, M. L. 1938. Estimation of pepsin, papain and cathepsin with hemoglobin. *J. Gen. Physiol.* 22: 79-89
- (16) Antweiler, H. J., and G. Klemmer. 1970. On the suitability of various methods of measurements for quantitative analysis of solutions of protein and proteide mixtures. *Zeitschrift fur Analytische Chemie.* 252(3/4): 256-259
- (17) Aoyagi, T., et al. 1969. Biological activities of leupeptins. *J. Antibiot.* 22(11): 558-568
- * * (18) Arnold, A., et al. 1951. Demonstration of the in vivo activity of papain. *Ann. New York Acad. Sci.* 54(2): 273-276
- (19) Arnon, R., and E. Shapira. 1967. Selective fractionation of antibodies against papain according inhibitory capacity. *Isr. J. Chem.* 5(4A): 105
- (20) Arnon, R., et al. 1969. Antibodies to papain. *Biochemistry* 6: 3942-3950

- (21) Arshad, M., et al. 1964. Preparation of an active meat tenderizer and its effect at various stages of cooking. Pakistan J. Sci. Res. 16(3/4): 151-156
- (22) Asatiani, V. S., and K. S. Mudzhiri. 1956. Papain. Sbornik Trudov Tbilissk. Nauch. - Issledovatel. Khim.-Farm. Inst. 8: 139-144
- (23) Austrian Patent 136 723. Pharmazeutische Werke "Norgine" A.-G. 1934. Steigerung der antianämischer Wirkung der tienschen Leber oder Niere.
- (24) Babin, R., et al. 1957. Testing a mixture of papain powder and bacitracin. Congr. soc. pharm. France, 9th Ed., Clermont-Ferrand. 1957, 117-121
- (25) Bagdy, D., et al. 1967. Effect of proteolytic enzymes on the activity of lipoprotein lipase in plasma. Folia Haematol. 87(4): 339-343
- (26) Bahadur, K., and I. Kumari. 1959. Fractionation of commercial papain by acetone and investigation of the proteolytic and milk clotting properties of the different fractions. Enzymologia 21: 114-122
- (27) Bahadur, K., and I. Kumari. 1959. Fractionation of papain by ethyl alcohol and investigation of the proteolytic and milk clotting properties of different enzymic fractions. Enzymologia 21: 145-154
- (28) Bahadur, K., and B. D. Atreya. 1960. Study of the effect of ripe papaya extract on the proteolytic activity of papain. Enzymologia 21(5): 325-336
- (29) Bahadur, K., and I. Kumari. 1960. Effect of heat on cupri-papain as indicated by its milk clotting and proteolytic activities. Enzymologia 21(6): 365-371
- (30) Baker, J. L., and H. F. E. Hulton. 1922. Amylasen der Getreidkörner. Die unlosliche Amylase der Gerste. J. Chem. Soc. London 121: 1929-1934

- (31) Balls, A. K., and W. S. Hale. 1935. Beeinflussung der Teigkonsiste. United States Patent 2,103,443
- (32) Balls, A. K., and S. R. Hoover. 1937. Die milchkoagulierende Wirkung von papain. J. Biol. Chemistry 121: 737-745
- (33) Balls, A. K., et al. 1937. Krystallisiertes Papain. Science 86: 379
- (34) Banga, I. 1952. The enzymic breakdown of the structure proteins of the aorta. Congr. intern. biochem. Resumes communs., 2^e Congr., Paris 1952, 269-270
- (35) Banner, R. E., et al. 1942. Action of certain enzymes, fish pancreatic islets and fetal calf pancreas on insulin. Endocrinology 31(2): 271-275
- (36) Baranova, V. Z., et al. 1960. Effect of deoxyribonucleic acid on enzymic action of papain and phosphoglyceric aldehyde dehydrogenase. Doklady Akad. Nauk S. S. S. R. 130: 450-452
- (37) Barta, G., and W. E. Parker. 1967. Papain activity in beer. Tech. Quart., Master Brew. Ass. Amer. 4(4): 254-256
- (38) Barton, A. W. 1938. Proteolytic vegetable ferment. Tr. Kansas Acad. Sc. 41: 165-166
- (39) Basu, K. P., and R. Chakravarty. 1935. Action of dyestuffs and narcotics on proteolytic enzymes. Trypsin and papain. J. Indian Chem. Soc. 12(2): 82-88
- (40) Baum, L. 1939. The local action of subcutaneous injections of enzymes. Frankfurt. Z. Path. 53: 220-233
- (41) Bazin, S., et al. 1962. Biochemical effects of crude papain on cartilaginous bone rudiments of chicken embryos. Ann. Inst. Pasteur 102(3): 346-352

- (42) Bednarski, W., et al. 1970. Utilization of some N compounds by moulds for the biosynthesis of proteins in a whey medium. *Przemysl Spozywczy* 24(11): 464-467
- * (43) Bender, M. L., et al. 1966. The determination of the concentration of hydrolic enzyme solutions -chymotrypsin, trypsin, papain, elastase, subtilisin and acetylcholinesterase. *J. Amer. Chem. Soc.* 88(24): 5890-5913
- (44) Bercovci, B., et al. 1959. Lytic property of human cord blood serum. *Proc. Soc. Exptl. Biol. and Med.* 100(4): 846-847
- (45) Berger, E. 1952. Activation of hemolysis and hemagglutination by papain. *Experientia* 8(6): 227-228
- (46) Bergmann, M., et al. 1937. Proteolytic enzymes. XV. Regarding the general nature of intracellular proteolytic enzymes. *J. Biol. Chem.* 119: 35-46
- (47) Bergmann, M., and C. Niemann. 1937. Proteolytic enzymes. XIV. The general nature of the enzymic degradation of proteins. *J. Biol. Chem.* 118: 781-788
- (48) Berkowitz-Hundert, R., et al. 1965. Milk clotting and proteolytic activity. *Enzymol Acta Biocatalytica* 29(2): 98-100.
- (49) Berkowitz-Hundert, R. 1966. Milk coagulation and proteolysis. *II Enzymologia* 31(5): 281-295
- (50) Bernfeld, P., and T. F. Kelley. 1964. Proteolysis of human serum β -lipoprotein. *J. Biol. Chem.* 239(10): 3341-3346
- (51) Bernheim, F. F. 1940. The action of papain and trypsin on certain dehydrogenases. *J. Biol. Chem.* 133(1): 141-144
- (52) Bersin, T., and H. Koster. 1935. Über den Einfluss von Oxydations-und Reduktions mitteln auf die Aktivitat von papain. *Hoppe-Seyler's Zeitschr. Physiol. Chem.* 233(1/2): 59-66

- (53) Bhatia, B. S., et al. 1967. Use of papain in the preparation of quick cooking dehydrated pulses and beans. Food Technol. 21(10): 105-107
- (54) Bhussry, B. R., et al. 1964. Wound healing in papain tested rats. Proc. Soc. Exp Biol Med. 115: 1010-1013
- (55) Binet, L., and D. Quivy. 1956. Mobilization of cutaneous histamine in the rat; action of various substances injected locally. Compt. rend. soc. biol. 150: 655-656
- (56) Binet, L., et al. 1956. Papain-induced shock in the rabbit. Compt. rend. soc. biol. 150: 1127-1131
- (57) Blair, G. W. S., and J. Burnett. 1958. Physical changes in milk caused by the action of rennet. I. Description of apparatus for measuring rigidity mouldli and internal viscosities, tests of reliability, and some observations on syneresis. J. Dairy Research. 25: 397-303
- (58) Blumberg, B. S., and A. G. Ogston. 1957. Effects of proteolytic enzyme on the hyaluronic acid complex of ox synovial fluid. Biochem. J. 66: 342-346
- * * (59) Blumenkrantz, N., et al. 1971. The effect of intravenous papain cartilage glycosaminoglycans. J. Invest Dermatol. 56: 381-383
- (60) Bohn, L. J., and C. H. Bailey. 1937. Wirkung der Gärung gewisser Teigbestandteile und Proteasen auf die physikalisch von Mehltegen. Cereal Chem. 14: 335-348
- (61) Bondi, A., and Y. Birk. 1954. Action of papain on protein feeds. Bull. Research Council Israel 4: 57-62
- * (62) Bonnet, R., and E. Meyer. 1951. Physiological synthesis of hemoglobin. IV. Role of papain in treatment of simple anemia. Bull. soc. chim. biol. 33: 1564-1570

- (63) Borchardt, W. 1928. Die Blausaureresistenz des d'Herelleschen Bakteriophagen. Ztschr. Immunitätsforsch. exp. Therapie 54: 403-408
- (64) Borroni, M., et al. 1963. Action of papain on the metaphysial growth zone. Arch. Ortop. 76: 513-520
- (65) Bose, S. M., and W. M. Krishna. 1957. Standardization and evaluation of different methods for the quantitative estimation of protease activity. Bull. Central Leather Research Inst., Madras 3: 329-342
- (66) Bosmann, H. B. 1968. Cellular control of macromolecular synthesis: Rates of synthesis of extracellular macromolecules during and after depletion by papain. Proc. Roy. Soc. Ser. B Biol. Sci. 169(1017) 399-425
- (67) Botsch, A. 1969. Possibilities for utilization of new enzymic baking additives. Brot. Gebaeck 23(10): 202-203
- (68) Bowlby, C., et al. 1953. A comparison of methods for determining proteolytic activity. Cereal Chem. 30: 480-491
- (69) Boyne, A. W., et al. 1967. Protein quality of feeding-stuffs: 4. Progress report on collaborative studies on the microbiological assay of available amino acids Brit. J. Nutr. 21(1): 181-206
- (70) Brading, I., et al. 1959. The effect of receptor-destroying enzyme, trypsin, and papain on red cell survival. J. Exptl. Biol. Med. Sci. 37: 37-48
- (71) Brill H. C., and R. E. Brown. 1922. Die Wirksamkeit von Philippinen-papain. Philippine J. of Science. 20: 185-191
- (72) British Patent. 1957. 782 809. Schering Corp.. Purified papains.

- (73) British Patent. 1959. 808 300. Schwarz Laboratories, Inc. Purification of papain
- (74) British Patent. 1969. 1 171 052. Hissink, Louis. Injectable antishock preparation.
- (75) British Patent. 1971. 1 255 867. Matsukura, K. Liquefied meat product.
- (76) Brocklehurst, K., and G. Little. 1970. Novel reactivity of papain and a convenient active site titration in the presence of other thiols. Fed. Eur. Biochem. Soc. Lett. 9(2): 113-116
- (77) Brooks, G. 1936. Reinigung von Abwassern kolloider Natur durch Papainase. Ann. Hyg. publ., ind. sociale. 14: 278-283
- (78) Brown, A., et al. 1952. The effect of trypsin and other enzymes on feline pneumonitis virus. J. Infectious Diseases 91: 184-190
- * * (79) Bryant, J. H., et al. 1958. The release of chondroitin sulfate from rabbit cartilage following the intravenous injection of crude papain. Arch. Biochem. and Biophys. 76(1): 122-130
- (80) Buffe, G. M. 1948. Proteolysis and lipophaneresis in blood serum. Annee biol. 52: 202-203
- (81) Bullock, K., and J. Kumar Sen. 1951. Papain assay process and standards. J. Pharm. Pharmacol 3: 476-485
- (82) Buluk, K., and A. Zuch. 1967. The possibility of proteolytic activation of the fibrin-stabilizing factor (FSF). Biochim. Biophys. Acta. 147(3): 593-594
- (83) Buluk, K., and A. Zuch. 1968. The similarity of activity between thrombin and papain in reaction with fibrinogen and fibrin stabilizers (FSF) Postepy Hig. Med. Dosw. 22(1): 177-188

- (84) Buschke, W. 1949. Studies on intercellular cohesion in corneal epithelium methods. Effects of proteolytic enzymes, salts, hydrogen-ion concentration, and nonpolar substances. J. Cellular Comp. Physiol. 33: 145-176
- (85) Calcinaï, M. 1933. Morphologic alterations induced in tissues by digestion with papain. Sperimentale 87(3/-4): 437-449
- (86) Calcinaï, M. 1939. Anaphylactogenic properties of the products of papain digestion of seroproteins and egg albumin. Boll. ist. sieroterap. Milan. 18: 594-613
- (87) Calcinaï, M. 1939. Commercial papain as an anaphylactogen. Boll. ist. sieroterap. Milan. 18: 677-682
- (88) Calf, G. E., and I. N. Jones. 1954. Composition for tenderizing meat. Australian Patent 157 454
- * (89) Capri, A., et al. 1961. Local and general action of basic proteins papain and trypsin. Boll. Ital. Biol. Sper. 37: 678-680
- (90) Carter, D. V., and G. Sykes. 1961. The assay of enzyme activity by the plate-diffusion technique. J. Pharm. Pharmacol. 13: Suppl. 195T-199T
- (91) Castaneda-Aguillo, M. 1956. Northrop's phenomenon. The digestion of proteins by enzymes in the presence of gelatin. J. Gen. Physiol. 39: 361-371
- (92) Castren, E. 1957. The determination of enzymic activity by the cup-plate method. Syensk Farm. Tidskr. 60: 871-878
- (93) Charavanapavan, C. 1945. Studies in the preparation properties and assay of commercial papain. Trop. Agric. 101(2): 72-76

- (94) Charlton, R. 1969. Seven pointers to a beef revolution. Farmers' Weekly 71(23): 59-62
- (95) Chatterjee, G., et al. 1957. Studies on the mechanism of blood coagulation by proteolytic enzymes. I. Effect of calcium on the clotting and proteolytic activities of papain and trypsin. II. Optimum pH for the proteolytic and clotting activities of papain and trypsin. Ann. Biochem. and Exptl. Med. India. 17(6): 179-186
- (96) Chatterjee, G., et al. 1958. Studies on the mechanism of blood coagulation by proteolytic enzymes. III. Role of inhibitors on the clotting and proteolytic activities of trypsin and papain. Ann. Biochem. and Exptl. Med. India. 18(1): 1-6
- (97) Chizhova, K. N. 1958. A study of gluten protein changes under the influence of separate factors related to the preparation of wheat dough. Trudy Vsesoyuz. Nauch-Issledovatel. Inst. Khlebopekar. Prom. 7: 112-120
- (98) Chrzaszcz T., and J. Janicki. 1934. Der Einfluß von Papain auf die amylolytische Kraft verschiedener Getreidearten. Biochem. Z. 272: 402-411
- (99) Chrzaszcz, T., und J. Janicki. 1934. Zusammenwirkung der proteolytischen Enzyme bei Erhöhung der Amylasemenge von verschiedenen Getreidearten. Roczniki Nauk rolniczych i lesnych 33: 191-200
- (100) Chrzaszcz, T., and J. Janicki. 1934. The presence of a kinase of amylase in trypsin preparations. Biochem. J. 28(6): 1949-1954
- * (101) Clagett, J. A., and W. E. Engelhard. 1967. Immunosuppressive activity of the 3a-1 fraction of papain. J. Bacteriol. 93(5): 1493-1498
- (102) Cole, E. R. 1968. Hydrolysis of L-histidine methyl ester. II. Activity of various proteolytic enzymes with special reference to activators of human plasminogen. Thromb. Diath. Haemorrh. 19(3-4): 334-345

- (103) Collier, B. 1966. Determination of papain in beer. J. Inst. Brewing. 72(2): 204-207
- (104) Copley, A. L., and V. Tsuluca. 1962. Inhibitory action of fibrinolysin and other proteolytic enzymes on capillary permeability induced by histamine, serotonin, and bradykinin. Life Sci. 1: 243-251
- (105) Corbin, H. L. 1956. Tenderization of meat. French Patent. 1 118 306
- (106) Corneli, W. 1931. Über die Einwirkung proteolytischer Fermente auf das Insulin. Ztschr. physiol. Chem. 199: 20-22/7.
- (107) Costa, A. L., et al. 1968. Effect in vitro of organic polycations on a pathogenic Candida albicans. Boll. Soc. Ital. Biol. Sper. 44(8): 731-735
- (108) Courtois, J. E. 1970. Determination of enzymes utilized in pharmacy. Therapeutique 46(1): 25-28
- (109) Crespo, J. G., and R. D. Salinas. 1964. Papain as a tenderizer of meats. Rev. Asoc. Bioquim. Agr. 29(155): 228-232
- (110) Crosti, P. F. 1961. Hematochemical modifications induced by intravenous administration of enzymes. I. Action of papain on the total proteins and on their electrophoretic fractions. Atti. Accad. Med. Lombarda 16: 141-143
- (111) Crosti, P. F., et al. 1961. Action of papain and other enzymes on connective tissue in vivo. Atti. Congr. Lega Intern. Reumatismo 10^o, Rome 2: 1008-1010
- (112) Crosti, P., et al. 1963. Serum proteins and glycoproteins in papain-injected rabbits. Am. J. Pathol. 43(3): 419-435

- (113) Davey, M. G., and E. F. Luescher. 1967. Actions of thrombin and other coagulant and proteolytic enzymes on blood platelets. *Nature* 216(5118) 857-858
- (114) Davidson, J., and J. A. LeClerc. 1942. Effect of various concentrations of papain and potassium iodate on the loaf volume of bread. *J. Agric. Res.* 64(3): 145-152
- (115) Davis, L. J., et al. 1943. Treatment of pernicious anemia with experimental proteolysed liver preparation; preliminary observations, *Brit. M. J.* 1: 655-656
- (116) Dawson, L. E., and G. H. Wells. 1969. Tenderness of freeze-dried chicken treated with proteolytic enzymes. *Poultry Science* 48(1): 64-70
- (117) Delaney, W. 1965. Tenderizing of fresh meats. *U. S. Patent* 3 188 213
- (118) Delaunay, A., et al. 1963. Lesions of the tracheal cartilage caused by injection of papain and their relation to the collagenoses observed in clinical medicine. *Minerva otorinolaring.* 13: 217-222
- (119) Delmotte-Plaquee, J. 1958. Assay methods for papain. *J. pharm. Belg.* 13: 227-242
- (120) Demaille, J., et al. Isolation of a glycopeptide fraction of α -macroglobulin in pig serum. *Soc. Chim. Biol. Bull.* 48(1) 45-52 1966
- (121) Denton, A. E., et al. 1966. Enzymic removal of meat from bones. *U.S. Patent* 3 293 687
- (122) Dhere, C., and C. Baumeler. 1929. Effects on hemocyanin of snail. *Compt. rend. Soc. de biol.* 101: 1071-1075
- (123) Dianzani, M. U. 1953. Action of papain and of trypsin on the morphology and some enzymatic activities of isolated mitochondria. *Experientia* 9(9): 343-345

- (124) Dorrington, K. J., et al. 1966. The proteolysis of immunoglobulin G with long-acting thyroid-stimulating activity. *Biochem. J.* 98(3) 853-861
- (125) Douste-Blazy, L., et al. 1962. Behavior of red cell lipids during in vitro hemolysis. *Biochem. Probl. Lipids, Proc. Intern. Conf., 7th, Birmingham, Engl.* 1962, 396-403
- (126) Dreyer, C., and P. C. Jones. 1969. Papain induced changes in cartilage. *J. Dent. Ass. S. Afr.* 24(2): 41-43
- * (127) Drezancic, I. I. 1957. Papain as an anthelmintic for *Haemonchus contortus* in sheep. *Australian Vet. J.* 33: 178-181
- (128) Dubouloz, P., and J. Laurent. 1953. A hematinlike pigment present in the liver of higher vertebrates. *Bull. soc. chim. biol.* 35: 781-786
- (129) Dyckerhoff, H., and D. Gigante. 1940. Occurrence of phytothrombin in papain; relation of papain to blood coagulation system. *Biochem. Ztsch.* 304: 334-339
- (130) Eagle, H., and T. N. Harris. 1937. Blood coagulation. V. The coagulation of blood by proteolytic enzymes (Trypsin, papain). *J. Gen. Physiol.* 20: 543-560
- (131) Edery, H. 1965. Further studies of the sensitization of smooth muscle to the action of plasma kinins by proteolytic enzymes. *Brit. J. Pharmacol.* 24(2): 485-496
- (132) Eisen, M. J., and A. Langer. 1938. Papain activators and cathepsin in tumors and the action of cysteine. *Enzymologia* 2: 321-328
- (133) Ellis, W. J., and F. G. Lennox. 1943. Some observations on Australian-grown papain. *Australia Counc. Sci. and Indust. Res. J.* 16(3): 166-172

- * * (134) Emele, J. F., et al. 1966. The analgesic-anti-inflammatory activity of papain. Arch. Int. Pharm. Codyn. Therap. 159(1): 126-134
- (135) Enders, C., et al. 1939. Determination of beer stability. II. Wochschr. Brau. 56: 329-332
- (136) Engfeldt, B., et al. 1961. Studies on the effect of papain on growing bone. Acta Path Microbiol. Scand. 51(Suppl 144): 43-45
- (137) Engfeldt, B. 1969. Studies on the epiphyseal growth zone; IV. The effect of papain on the ultrastructure of epiphyseal cartilage. Acta Pathol. Microbiol. Scand. 75(2): 220-228
- (138) Erbring, H., and F. Corcillius. 1951. Measurements of enzymic activity of the juice of Carica papaya. Deut. Apoth-Ztg. 91: 91-93
- (139) Erdos, J. 1950. A method for the determination of the activity of papain. Enzymologia 14: 8-9
- (140) Erdos, J., and G. Cortes. 1957. Hydrolysis assay of adrenal gland. Ciencia. 17: 27-30
- (141) Erlandson, A. L. Jr. 1966. The induction of non-specific resistance in mice by papain. J. Infect Dis. 116(3): 297-302
- (142) Erlich V. L., and G. M. Burkert. 1949. Influence of salts on amylase of malt. Cereal Chem. 26: 239-257
- * * (143) Ershoff, B. H., et al. 1963. Submaxillary gland hypertrophy in rats fed proteolytic enzymes Proc. Soc. Exp. Biol. Med. 113: 879-881

- (144) Esh, G. C., and J. M. Som. 1958. Enzymic activity of papain. I. Peanut cake as the substrate. J. Proc. Inst. Chemists 30: 191-196
- (145) Everett, A. L. 1960. In situ histological evaluation of elastase activity. Stain Technol. 34: 325-330
- (146) Eyk, H. G., et al. 1967. Fragmentation of human IgG globulin with papain trypsin and pepsin. Clin. Chim. Acta 16: 429-431
- (147) Eyquem, A., and Y. Crepin. 1956. Appearance of autoagglutinin of high titer in rabbits or dogs immunized with papain treated red blood cells. Ann. Inst. Pasteur 90: 364-367
- (148) Faillard, H., et al. 1962. Action of neuraminidase and papain on intrinsic factoractive mucoids from the stomachs of men and swine. Z. Physiol. Chem. 327: 100-108
- (149) Falk, K. G. 1933. Richtunggebende Einflüsse in biologischen Systemen. III, Der Einfluß von Proteasen auf Lipasewirkungen. J. biol. Chemistry 103: 363-372
- (150) Farrell, J. H. 1956. Effect on digestibility of methods commonly used to increase the tenderness of lean meat. Brit. J. Nutrition 10: 111-115
- (151) Fausel, E. G. 1934. Experimental postoperative intra-abdominal adhesions with special reference to use of papain solution. M. Times & Long Island M. J. 62: 1-4
- (152) Fauve, R. M., and C. H. Pierce-Chase. 1967. Comparative effects of corticosteroids on host resistance of infection in relation to chemical structure. J. Exp. Med. 125(5): 807-821
- * (153) Feinberg, H., et al. 1952. Antifatty liver action of papain and ficin in insulin-treated, depancreatized dogs. Proc. Soc. Exptl. Biol. Med. 80: 161-162

- (154) Fell, H. B., and L. Thomas. 1960. Comparison of the effects of papain and vitamin A on cartilage. II. The effects on organ cultures of embryonic skeletal tissue. J. Exp. Med. 111: 719-744
- (155) Fels, G., et al. 1952. Enzymatic digestion of beef thyrotropic hormone with papain. Endocrinology 51(5): 349-352
- (156) Felton, F. D., and G. Kauffmann. 1927. Enzymwirkung auf Pneumokokkenantikörper. J. Immunology 13: 219-235
- (157) Ferguson, J. H., and P. H. Ralph. 1943. Ninhydrin, crystalline papain and fibrin clot. Am. J. Physiol. 138: 648-651
- (158) Follis Jr., R. H. 1959. Spontaneous hemorrhage induced by administration of aminoacetonitrile and papain of anticoagulants to rats. Proc. Soc. Exptl. Biol. Med. 102: 114-118
- * (158a) Food Chemicals Codex. 1972. 2nd. Ed. National Academy of Sciences. Washington, D. C. p. 574
- (159) Ford, W. P., and A. M. Maiden. 1938. Die im Teig durch Glutathion und Papain hervorgerufenen Effekte. J. Soc. chem. Ind. 57: 278-281
- (160) Forstner, G. G. 1971. Release of intestinal surface-membrane glycoproteins associated with enzyme activity by brief digestion with papain. Biochem. J. 121: 781-789
- * * (161) Fosdick, L. S., and G. W. Rapp. 1944. Effect of proteolytic enzymes on acid formation in the mouth. J. Dental Research 23: 81-83
- (162) Frankel, E. M. 1921. Studien über Enzymwirkung. XV. Faktoren, welche die proteolytische Wirksamkeit des papains beeinflussen. J. Biol. Chem. 31: 201-215

- (163) Frassi, G. A., et al. 1963. Evolutive aspects of the changes of the growth zone caused by papain. Arch. Ortop. 76: 493-502
- (164) French Patent. 1932. 733 241. Wallerstein Co., Inc. Herstellung von Schokoladensirup.
- (165) French Patent. 1934. 763 482. Lunt, W. R. F. Behandlung von Nahrungsmitteln.
- (166) French Patent. 1966. 1 455 265. Swift & Co. Treatment of meat.
- (167) French Patent. 1967. 1 494 282. Ponomarett, T. Meat tenderizer containing papain stabilized with dextrose.
- (168) French Patent. 1969. 1 576 222. Hawley, R. L. Cooked egg preparation.
- (169) Freudenberg, K., et al. 1932. Die Wirkung proteolytischer Fermente auf Insulin und seine Dervate. XII. Mitt. uber Insulin. Hoppe-Seyler's Z. physiol. Chem. 213: 248-262
- (170) Frey, E. K., and H. Kraut. 1928. Nachweis und Wirkung eines Kreislaufhormons. Münch. med. Wchschr. 75: 763-764
- (171) Froissard, R. 1926. Das Papain des Handels. Seine Reinigung. Bull. Soc. Chim. Biol. 8: 288-293
- (172) Frossard R., and R. Fabre. 1925. Einfluß der Reaktion des Mediums auf die Verdauung durch Papain. C. R. Soc Biol. 92: 59-60
- * *(172a) Furia, T. E. 1968. Handbook of Food Additives. The Chemical Rubber Co. Press. Cleveland, Ohio. pp. 52, 67, 87

- (173) Gaetani, E. 1946. Effects of carcinogenic hydrocarbons on enzymatic functions of papain. *Tumori* 20: 165-170
- (174) Gauthier, B., et al. 1957. Assay of Papain. *Ann. Pharm. franc.* 15: 31-39
- (175) Gawron, O., and F. Draus. 1956. Modified milk clotting procedure for determination of papain activity. *J. Agr. Food Chem.* 4: 349-352
- (176) Geoffroy, R., and G. Labour. 1934. Wirkung der proteolytischen Enzyme auf die Proteine der Weizenmehle. *Bull. Soc. Chim. biol.* 16: 1625-1630
- (177) Gergely, J., and D. R. Stanworth. 1966. Papain-sensitive and papain-insensitive populations of IgG. *Haematol. Hung.* 6(1-2) 19-27
- (178) Gergely, J., et al. 1967. Structural studies of immunoglobulins. I. The role of cysteine in papain hydrolysis. *Immunochemistry* 4(2): 101-111
- (179) Gergely, J., et al. 1967. Structural studies of immunoglobulins. II. The varying susceptibility to papain-digestion of a group of human myeloma γ G-globulins. *Immunochemistry* 4(6): 369-374
- (180) German Patent. 1954. 903502. Koch, A., et al. Growth-promoting substance.
- (181) German Patent. 1960. 1 095 093. Mohler, H. Preservation of Meat.
- (182) Girard, H. 1962. The action of papain on the embryonic tibia of fowl cultivated in vitro. *J. Embryol. and Exptl. Morphol.* 10(3): 231-247
- (183) Gol'dshtein, B. I., and A. S. Tsiperovich. 1940. The synthetic action of papain. *Biochem. J. (Ukraine)* 16: 165-192

- (184) Gortner, W. A. 1954. Effect of age on enzymic release of amino acids from aorta tissue. J. Gerontol. 9(3): 251-261
- (185) Gotterer, G. S., and H. G. Williams-Ashman. 1957. Some factors which influence vesiculase action. Proc. Soc. Exptl. Biol. Med. 94: 60-64
- (186) Gottschall, G. Y., and M. W. Kies. 1942. Digestion of beef by papain. Food Res. 2(5): 373-381
- * * (187) Grant, R. A., et al. 1960. Chemical changes in rabbit aorta and ear cartilage induced by papain and calciferol. Biochem. J. 76: 412-415
- (188) Grassmann, W. 1935. Zur Kenntnis des natürlichen Papainaktivators. Biochem. Z. 279: 131-136
- (189) Greenawald, K. A., and T. T. Tsaltas. 1964. Papain induced chemical changes in adult rabbit cartilage matrix. Proc. Soc. Exp. Biol. Med. 117(3): 885-888
- (190) Greenway, P. J., et al. 1953. The papaw- its botany, cultivation, diseases, and chemistry. Tanganyika Territ. Dept. Agr. Pamphlets. No. 52, 32 pp.
- (191) Grieco, F. 1934. Prevention of postoperative peritoneal adhesions; experimental research. Ann. Ital. di chir. 13: 269-286
- (192) Grimes, O. F., et al. 1954. Effects, clinical report on cytologic diagnosis of gastric cancer. Surg. Gynec. & Obst. 98: 347-352
- (193) Griswold, R. M. 1955. Effect of different methods of cooking beef round of commercial and prime grades. I. Palatability and shear values. Food Research 20: 160-170

- (194) Groll, H. 1937. Über die Einwirkung von Fermenten auf Gewebsschnitte. Zeitschr. Ges. Exp. Med. 101(4): 451-456
- (195) Groll, H. 1926. Experimentelle Untersuchungen zur Lehre von der Entzündung. Krankheitsforsch. 2: 220-262
- (196) Grossgebauer, K. 1967. Biological activities of polyelectrolytically active macromolecules. Arch. Hyg. Bakteriол. 151(3/4) 360-370
- (197) Gualtieri, G., et al. 1964. Alterations of articular cartilage due to papain. Arch. Ortop. 77: 557-565
- (198) Gulland, J. M., and T. F. Macrae. 1933. Wirkung von proteolytischen Enzymen auf das oxytocische Prinzip der Hypophyse. Nature (London) 131: 470
- (199) Guri, C. D., and D. S. Bernstein. 1967. Rat epiphyseal cartilage: IV. Effect of papain on proteinpolysaccharide metabolism and S³⁵ incorporation, in vitro and in vivo. Proc. Soc. Exp. Biol. Med. 124(2): 380-386
- (200) Guven, C., and G. Ozsari. 1967. Identification of lipase, diastase, papain, emulsin, and invertase by thin-layer chromatography. Eczacilik Bul. 9(1): 12-14
- (201) Haas, R. 1940. Antigenic properties of papain preparations. Biochem. Z. 305: 280-289
- (202) Hadidian, Z., and M. M. Murphy. 1955. The interactions of the nonspecific inhibitor of hyaluronidase with hyaluronidase and proteolytic enzymes in vivo. J. Gen. Physiol. 39: 185-195
- (203) Hakim, A. A., and R. L. Peters. 1962. Enzyme specific action in vivo. I. Action of certain enzymes by the intravenous route. Exptl. Med. Surg. 20: 194-211

- (204) Hale, M. B. 1969. Relative activities of commercially-available enzymes in the hydrolysis of fish protein. Food Technology 23(1): 107-110
- (205) Halliwell, G. 1954. Action of proteolytic enzymes on Clostridium parbotulinum toxin. Biochem. J. 58: 4-8
- (206) Halpern, B. N., et al. 1963. Induction in rabbits of tissue lesions of autoimmune type and demonstration of anti-tissue antibodies after prolonged treatment with papain. Compt. Rend. Soc. Biol. 157: 225-231
- (207) Halpern, B., et al. 1964. Osteo-cartilaginous dystrophy and nanism experimentally induced in the monkey and rabbit by papain administration. C. R. Acad. Sci. 258: 6583-6588
- (208) Halpern, B. N., et al. 1965. Experimental collagen-like disease induced by repeated injections of papain, with evidence of autoimmune antibodies. Ann. N. Y. Acad. Sci. 124: 395-411
- (209) Halpern, B., et al. 1966. Osteo-cartilaginous dystrophies and dwarfism induced by papain in rabbits. Microradiographic and histochemical study. Rev. Franc. Clin. Biol. 11: 357-372
- (210) Hartiala, K., and M. I. Grossman. 1952. Studies on chemical and physical changes in duodenal mucus. J. Biol. Chem. 195(1) 251-256
- (211) Haurowitz, F. 1931. Splitting of hemoglobin. Ztschr. f. physiol.Chem. 194: 98-106
- (212) Haurowitz, F., et al. 1943. Digestibility of native and denatured egg albumin. Tib. Fakültesi (Istanbul) 6: 3231-3233
- (213) Haustein, K. O., and F. Markwardt. 1966. Effect of thrombin and other proteolytic enzymes on smooth muscle. Arch. Intern. Pharmacodyn. Ther. 163(2): 393-402

- (214) Hecht, A. 1967. Enzymichistochemical demonstration of different damage mechanisms in rat caridac muscles; phosphorylase activity. Beitr. Pathol. Anat. Allg. Pathol. 135(3): 322-332
- (215) Hecht, A. 1967. Enzyme-histochemical response of the liver to intravenous administration of papain. Acta. Biol. Med. Ger. 18(3): 397-407
- (216) Hedberg, M., and J. K. Miller. 1964. The effect of micro-organisms in blood. N. Y. State Dept. Health, Ann. Rept. Div. Lab. Res. 1964: 114
- (217) Heise, E., and M. Goerlich. 1963. Proteolytic activation of glucose-6-phosphatase in normal, diabetic, and hepatomic rat livers. Nature 197: 1311-1312
- (218) Helm, E. 1936. Biertrübungen aus nichtbiologischer Ursache. J. Inst. Brewing 42: 379-393
- (219) Hidaka, Y. 1960. The inhibitory effect of vegetable and bacterial proteolytic enzymes upon the growth of Ehrlich ascites cancer in mice. Toho Igakkai Zasshi. 7: 969-978
- (220) Hinkel, E. T. Jr., and C. E. Alford. 1951. Methods for measuring the proteolytic activity of papain. Ann. N. Y. Acad. Sci. 54: 208-227
- * * (221) Hinkel, E. T. Jr., and A. Arnold. 1951. The effect of the activity state of the enzymes on the in vivo activity of papain. Ann. N. Y. Acad. Sci. 54(2): 277-281
- (222) Hites, B. D., et al. 1951. Study of proteolytic activity in wheat flour doughs and suspensions. II. A papain inhibitor in flour. Cereal Chem. 28: 1-18
- (223) Hochwald, G. M., and G. J. Thorbecke. 1963. Trace proteins in cerebrospinal fluid and other biological fluids. II. Effect of storage and enzymes on the electrophoretic mobility of γ -trace and β -trace proteins in cerebrospinal fluid. Clin. Chim. Acta 8(5): 678-684

- (224) Hohmann, W. 1954. Significance of the pretreatment of the point of inoculation-for the take and growth of the mouse ascites carcinoma. Z. Krebsforsch. 59: 640-651
- * * (225) Holsinger, J. W. Jr., et al. 1968. Esophageal perforation following meat impaction and papain ingestion. J. Amer. Med. Ass. 204(8): 734-735
- (226) Hoover, S. R., and E. L. C. Kokes. 1947. The effect of pH upon proteolysis by papain. J. Biol. Chem. 167(1): 199-207
- (227) Horie, Y. 1961. Protein-clotting activity of papain. Agric. and Biol. Chem. 25(12): A97
- (228) Horie, Y. 1961. Microassay of papain and other proteases. Nippon Nogeikagaku Kaishi 35: 1361-1364
- (229) Horie, Y. 1962. A method for the quantitative estimation of papain in beer. Rept. Res. Lab., Kirin Brewery Co. Ltd. 5: 25-32
- (230) Horie, Y. 1962. Casein-, egg-albumin-and hemoglobin-clotting activities of papain. Agric. and Biol. Chem. 26(3): A(18-19):
- (231) Horie, Y. 1964. Protein-clotting activity of papain. Am. Soc. Brewing Chemists 1964, 174-182
- (232) Horie, Y. 1965. Electrophoretic studies of the interaction between protease and RNA. Rept. Res. Lab. Kirin Brew. Co., Ltd. No. 8: 7-11
- (233) Hovig, T. 1965. Effect of various enzymes on the ultrastructure, aggregation and clot-retraction ability of rabbit blood platelets. Thromb., Diath. Haemorrhag. 13(1): 84-113

- (234) Howell, D. S., and L. Carlson. 1965. The effect of papain on mineral deposition in the healing of rachitic epiphyses. *Exptl. Cell Res.* 37(3): 582-596
- (235) Hoyt, R. E., and H. Zwicker. 1952. The effect of human serum upon the agglutination of enzyme-treated erythrocytes. *J. Immunol.* 68(1): 61-64
- (236) Hoyt, R. E., and H. Zwicker. 1953. The role of enzyme in reversible agglutination of red cells. *J. Immunol.* 71: 325-330
- (237) Hsiao, S., and F. W. Putnam. 1961. Cleavage of human γ -globulin by papain. *J. Biol. Chem.* 236: 122-135
- (238) Huet, R. 1956. Nature and properties of papain. *Fruits* 11: 51-57
- (239) Huffman, D. L., et al. 1961. The effect of antemortem injection of papain on tenderness of chicken. *Poultry Sci.* 40(6): 1627-1630
- (240) Huffman, D. L., et al. 1967. Effect of antemortem injection of sodium chloride, papain, and papain derivatives on the tenderness of beef. *J. Anim. Sci.* 26(2): 285-289
- (241) Huffman, D. L., et al. 1967. Effect of breeding level of feeding and antemortem injection of papain on the tenderness of weanling calves. *J. Anim. Sci.* 26(2): 290-293
- (242) Hulth, A. 1958. Experimental retardation of endochondral growth by papain. *Acta. Orthopaed. Scan.* 28(1): 1-21
- (243) Hulth, A. 1958. The growth inhibiting effect by papain on young rabbits. *Acta. Orthopaed. Scan.* 27(3): 167-172
- (244) Hulth, A., and O. Westerborn. 1959. Experimental production of dwarfs in rabbits with papain. *Exptl. Cell Research.* 17: 543-547

- (245) Hummel, K. 1965. Comparison of the action of cholera neuraminidase with other plant and animal enzymes liberating neuraminic acid, on the physicochemical properties of human erythrocytes. *Pathol. Microbiol.* 28(6): 891-900
- (245a) Hurst, T. L., and S. W. Fox. 1956. The course of proteolysis of lysozyme. *Arch. Biochem and Biophys.* 163(2): 352-367
- (246) Hutchins, B. K., et al. 1967. Effects of additives and refrigeration on reducing activity, metmyoglobin, and malonaldehyde of raw ground beef. *J. Food Sci.* 32(2): 214-217
- (247) Hutchinson, C. 1964. Effects at a single metaphysis of papain. *Anat. Rec.* 148(2): 293
- (248) Ichishima, E. 1967. Structure and activity of proteolytic enzymes. *Tampakushitsu, Kakusan, Koso* 12(7): 539-552
- (249) Ilany-Feigenbaum, J. 1966. Milk coagulation and proteolysis I. *Enzymologia* 31(5): 274-280
- (250) Indian Patent. 1971. 124577. Hindustan Lever Ltd. Dried Milk Products
- (251) Indian Patent. 1971. 124578. Hindustan Lever Ltd. Dried Milk Products
- * (252) Innerfield, I., and T. Wernick 1961. Plasma anti-thrombin alterations following oral papain. *Proc. Soc. Exptl. and Med.* 107(3): 506-507
- * (253) Innerfield, I., et al. 1963. Peptidase activity in rat urine following orally given proteases. *Life Sci.* 2(9): 671-675

- * (254) Innerfield, I., et al. 1964. Tissue peptidase activity following orally given proteases. Life Sci. 3(3): 267-275
- * (255) Innerfield, I., et al. 1965. "Fibrin-to-clot" clotting time after oral streptokinase or papain. Life Sci. 4(13): 1301-1305
- (256) Irimajiri, S., et al. 1968. Some chemical and biological properties of the F'c fragment of human γ G globulins. Immunochemistry 5(4): 383-397
- (257) Irving, J. T. 1960. The effect of papain on the staining properties of bone and teeth. Arch. Oral Biol. 2: 241-242
- (258) Irving, J. T., and O. V. Ronning. 1962. The selective action of papain on calcification sites. Arch. Oral Biol. 7(5): 357-363
- (259) Italian Patent. 1955. 532.410. Moris, Maurizio. Mixtures for the solubilization of proteins and particularly for the rapid disintegration of the fibers of edible meats.
- (260) Itoh, K. 1966. Comparative studies on the protease inhibitors in normal or inflammatory serum of rabbit. Mie Med. J. 16(1): 9-24
- (261) Jacoby M. 1926. Zur Kenntniss der Bedeutung der Blausäure bei der Metallvergiftung der Enzyme. Biochem. Ztschr. 175: 79-85
- (262) Jacquet, J., et al. 1968. Coagulation of milk by enzymes. C. R. Hebd. Seances Acad. Agr. Fr. 54(4): 254-264
- (263) Japanese Patent. 1946. 174192. Ishida, N. Nutrient containing much yeast and glycogen.

- (264) Japanese Patent. 1970. 9212 70. Yasukawa, H.
Liquefied egg product.
- (265) Jefferis, R., et al. 1968. Relation between the
papain sensitivity of human γ G-immunoglobulins
and their heavy chain subclass. Nature 219(5154):
646-649
- (266) Jirasek, J. E. 1963. Effect of papain on various
tissues. Sborn,Ved,Prac,Lek,Fak,Karlov Univ, 6:
Suppl: 335
- (267) Johansen, J. T., and M. Ottesen. 1968. Proteolytic
degradation of the B-chain of oxidized insulin by
papain, chymopapain and papaya peptidase. C. R.
Trav. Lab. Carlsberg 36(15): 265-283
- * * (268) Johanson, W. G. Jr., et al. 1971. The evolution of
papain emphysema in the rat. J. Lab. Clin.
Med. 78: 599-607
- (269) Johanson, W. G., et al. 1972. Effects of elastase
collagenase, and papain on structure and function
of rat lungs in vitro. Clin. Invest. 51:
288-293
- (270) Johnson, J. A., et al. 1956. Properties of certain
protease systems used in breadmaking. Cereal
Chem. 33: 1-17
- (271) Johnston, B. 1956. Thiolsterase activity in papain.
J. Biol. Chem. 221: 1037-1046
- (272) Johnston, W. R., and S. Jozsa. 1935. Eine allgemeine
Methode zur Bestimmung der Konzentration von
Enzympräparaten. J. Amer. Chem. Soc. 57:
701-706
- (273) Jones, M., et al. 1967. Peptidase activity in wort
and beer. Amer. Soc. Brew. Chem., Proc., 1967,
14-17

- (274) Joseph, R. L. 1970. Production of tender beef. Process Biochemistry 5(11): 55-58
- (275) Jozsa, S., and H. C. Gore. 1932. Development of diastatic enzymes of malt and flour. Effect of certain salts carbohydrates, and papain. Ind. & Engineer. Chem. 24(1): 95-98
- * * (276) Jozsa, L., and G. Szederkenyi. 1968. Enzymic dissolution of kidney stones. Z. Urol. Nephrol. 61(5): 295-300
- (277) Kaemmerer, K., and L. V. Erichsen. 1954. The use of preparations of plant enzymes. Monatsh. Tierheilkunde 6: 128-136
- (278) Kaleel-Head M. R. 1964. Changes in certain characteristics of bovine muscle when treated with papain, hyaluronidase and lysozyme. Dissertation Abstr. 24(12): 5375
- (279) Kalliomaki, J. L., et al. 1961. An attempt to provoke rheumatoid factor-like phenomenon in rabbit serum by repeated intravenous injections of papain. Ann. Med. Exp. Fenn. 39: 62-65
- (280) Kalmanson, G. M., and J. Bronfenbrenner. 1942. The reversal of Pneumococcus Quellung by digestion of the antibody with papain. Science 96(2479): 21-22
- (281) Kalmanson, G. M., and J. Bronfenbrenner. 1942. The reactivation of neutralized phage and other biologic agents by digestion of the antibody with papain. Federation Proc. Pt. II 1(1): 179-180
- (282) Kalmanson, G. M., and J. Bronfenbrenner. 1943. Restoration of activity of neutralized biologic agents by removal of antibody with papain. J. Immunol. 47: 387-407
- (283) Kambara, T., et al. 1964. Induction of mitoses in the livers of rats by means of intraperitoneal papain. Kumamoto Med. J. 17(1): 18-30

- (274) Joseph, R. L. 1970. Production of tender beef. *Process Biochemistry* 5(11): 55-58
- (275) Jozsa, S., and H. C. Gore. 1932. Development of diastatic enzymes of malt and flour. Effect of certain salts carbohydrates, and papain. *Ind. & Engineer. Chem.* 24(1): 95-98
- * * (276) Jozsa, L., and G. Szederkenyi. 1968. Enzymic dissolution of kidney stones. *Z. Urol. Nephrol.* 61(5): 295-300
- (277) Kaemmerer, K., and L. V. Erichsen. 1954. The use of preparations of plant enzymes. *Monatsh. Tierheilkunde* 6: 128-136
- (278) Kaleel-Head M. R. 1964. Changes in certain characteristics of bovine muscle when treated with papain, hyaluronidase and lysozyme. *Dissertation Abstr.* 24(12): 5375
- (279) Kalliomaki, J. L., et al. 1961. An attempt to provoke rheumatoid factor-like phenomenon in rabbit serum by repeated intravenous injections of papain. *Ann. Med. Exp. Fenn.* 39: 62-65
- (280) Kalmanson, G. M., and J. Bronfenbrenner. 1942. The reversal of *Pneumococcus* Quellung by digestion of the antibody with papain. *Science* 96(2479): 21-22
- (281) Kalmanson, G. M., and J. Bronfenbrenner. 1942. The reactivation of neutralized phage and other biologic agents by digestion of the antibody with papain. *Federation Proc. Pt.II* 1(1): 179-180
- (282) Kalmanson, G. M., and J. Bronfenbrenner. 1943. Restoration of activity of neutralized biologic agents by removal of antibody with papain. *J. Immunol.* 47: 387-407
- (283) Kambara, T., et al. 1964. Induction of mitoses in the livers of rats by means of intraperitoneal papain. *Kumamoto Med. J.* 17(1): 18-30

- * (284) Kambara, T., and Y. Nohara. 1966. Mitotic changes in rat liver after intraperitoneal papain protease. Arch. Pathol. 81(6): 525-530
- (285) Kang, C. K., and E. E. Rice. 1970. Degradation of various meat fractions by tenderizing enzymes. J. of Food Science 35(5): 563-565
- (286) Kaufmann, B., and S. P. Masouredis. 1963. Effect of formalin and periodic acid on red cell Rh. (D) and enzyme-exposed Rh. (D) antigen. J. Immunol. 91(2): 233-242
- (287) Kellner, A., and T. Robertson. 1954. Selective necrosis of cardiac and skeletal muscle induced experimentally by means of proteolytic enzyme solutions given intravenously. J. Exptl. Med. 99: 387-403
- * * (288) Kilburn, K. H., et al. 1971. Morphological biochemical assessment of papain-induced emphysema. Arch. Intern Med. 127: 884-890
- (289) Kim, Y. B., et al. 1964. Inactivation of gram-negative bacterial endotoxins by papain. Proc. Soc. Exp. Biol. Med. 115: 140-142
- (290) Kimmel, J. R., and E. L. Smith. 1954. Crystalline papain I. Preparation, specificity and activation. J. Biol. Chem. 207(2): 515-531
- (291) Kimmel, J. R., and E. L. Smith. 1958. Crystalline papain and benzoyl-L-argininamide. Biochem. Preparations 6: 61-67
- (292) Kind, L. S., and J. W. Goodman. 1962. Activity of pepsin and papain digested gamma globulin in mouse anaphylaxis. In: 46th Annual meeting, Atlantic City, New Jersey, April 1962. Fed. Proc. 21(2): 272
- (293) Kirk, J. S., and J. B. Sumner. 1934. The reaction between crystalline urease and antiurease. J. Immunol. 26(6): 495-504

- (293a) Kirk - Othmer Encyclopedia of Chemical Technology. 1967. 2nd. Ed. Volume 8. Interscience Publishers, New York. pp.188, 205.
- (294) Kitagawa, K., and N. Izumiya. 1959. Action of trypsin and papain on the derivatives of α -amino- γ -guanidinobutyric acid and homoarginine. J. Biochem. (Tokyo) 46: 1159-1165
- (295) Kitulescu, I., and V. Baluta. 1963. Regulation of the activity of proteases by physiological factors. Acad. Rep. Populare Romine Studii Cercetari Biochim. 6(4): 523-528
- (296) Kitulescu, I., et al. 1967. The use of papain and of mediators of the nervous activity: histamine and acetylcholine for differentiating glycoproteins in blood of patients with cancer. Rum. Med. Rev. 21: 70-73
- (297) Klemm, G., and W. Altermann. 1937. The influence of proteolytic enzymes on the gluten quality and baking properties of wheat flours. Z. ges. Getreide-, Muhlen-u. Backereiw. 24: 28-33
- (298) Kohn, R. R. 1962. Age-related variation in susceptibility of human muscle protein to enzymic digestion. Exptl. Cell Res. 26: 223-225
- (299) Koj, A., and A. C. Allison. 1969. Effects of agents labilizing lysosomes on concentrations of "acute-phase" proteins in rabbit plasma. Folia Biol. (Warsaw) 17(1): 37-47
- (300) Kono, T. 1969. Destruction of insulin effector system of adipose tissue cells by proteolytic enzymes. J. Biol. Chem. 244(7): 1772-1778
- (301) Kopaczewski, W. 1949. The action of some coagulants on the blood. Compt. rend. 229: 630-632
- (302) Kourilsky, R., et al. 1956. The immune-adhesion of the staphylococcus. Evidence of a specific antigen-antibody system. Ann. n st. Pasteur 90: 1-7

- (303) Kraut, H., and E. Bauer. 1927. Estimation, purification, chemical properties. Ztschr. f. physiol. Chem. 164: 10-36
- (304) Krebs, A. 1930. Proteolytic action. Biochem. Ztschr. 220: 289-303
- * * (305) Kyriazis, A. P. 1971. Studies in permeability of articular cartilage in New Zealand albino rabbits. The effect of aging, papain, and certain steroid hormones. Am. J. Pathol. 62: 75-85
- (306) Kulberg, A. Y., and I. A. Tarkhanova. 1961. Immunological study of papain-fermented antibodies. J. Hyg. Epidemiol. Microbiol. and Immunol. 5(4): 444-453
- (307) Kul'berg A. YA. et al. 1962. Immunological properties of papain-digested antibacterial and antirickettsial rabbit sera. Byull. Eksptl. Biol. i Med. 52(12): 1406-1408
- (308) Kul'berg, A. Ya. et al. 1966. Studies of serological properties of papain treated rabbit antibodies towards bacterial antigens of various chemical nature. Vop, Med, Khim. 12(5): 461-467
- (309) Kuo, J. F., et al. 1966. The effect of proteolytic enzymes on isolated adipose cells. Life Sci. 5: 2257-2262
- (310) Kuroda, K., and M. Fujino. 1954. Action of the proteolytic enzymes on heat-coagulated egg yolk. Tokushima J. Exptl. Med. 1: 85-91
- (311) Kuroiwa, Y., and H. Hashimoto. 1960. Fractionation of degraded products of lupulones and humulones by chromatographic method. Rept. Research Labs. Kirin Brewery Co. No. 3: 11-16.
- (312) Lack, C. H. 1962. Increased vascular permeability, chondrolysis and cortisone. Proc. Roy. Soc. Med. 55: 113-116

- (313) Laforte, M., et al. 1968. Effect of phenothiazine on the red cells of paroxysmal nocturnal hemoglobinuria. Brit. J. Haematol. 14(4): 345-350 .
- (314) Lamm, M., et al. 1968. Modulation of TL(thymus-leukemia) antigens by Fab-fragments of TL antibody. J. Immunol 101(1): 99-103
- (315) Lauwers, A. 1963. Standardization of proteinase and the comparative determination of papain. Arch. Int. Physiol. 71: 637-638
- (316) Lauwers, A. R. 1964. The standardization of proteinases. West European Symp. Clin. Chem. 4: 19-31
- (317) Lauwers, A., and R. Ruyssen. 1965. Determination of the activity of papain. J. Pharm. Belg. 20(1-2): 11-50
- (318) Lazo-wasem, E. A. 1966. Standardization of papain activity. Report of a collaborative study. J. Pharm. Sci. 55: 723-725
- (319) Lee, C. L., et al. 1964. Effect of papain on antigenicity of mouse erythrocytes. In: 48th Annual meeting of the Federation of American Societies for Experimental Biology, Chicago, 1964. Fed. Proc. 23 (2Pt.1): 389
- (320) Leipert, T., and I. Häfner. 1930. Über die Abspaltung von Tyrosin und Tryptophan aus Casein durch mit Blausäure aktiviertes Papain. Biochem. Ztschr. 229: 427-432
- (321) Leont'eva -Tyukina, A. A. 1967. Elastase activity of papain. Prof. Travmatizma, Lecheniya Travm Zabol. Oporno-Dvigat. App. 337-341
- (322) Leslie, R. G., et al. 1971. The products from papain and pepsin hydrolyses of guinea-pig immunoglobulins gamma 1G and gamma 2G. Biochem. J. 121: 829-837

- (313) Laforte, M., et al. 1968. Effect of phenothiazine on the red cells of paroxysmal nocturnal hemoglobinuria. Brit. J. Haematol. 14(4): 345-350
- (314) Lamm, M., et al. 1968. Modulation of TL(thymus-leukemia) antigens by Fab-fragments of TL antibody. J. Immunol 101(1): 99-103
- (315) Lauwers, A. 1963. Standardization of proteinase and the comparative determination of papain. Arch. Int. Physiol. 71: 637-638
- (316) Lauwers, A. R. 1964. The standardization of proteinases. West European Symp. Clin. Chem. 4: 19-31
- (317) Lauwers, A., and R. Ruysen. 1965. Determination of the activity of papain. J. Pharm. Belg. 20(1-2): 11-50
- (318) Lazo-wasem, E. A. 1966. Standardization of papain activity. Report of a collaborative study. J. Pharm. Sci. 55: 723-725
- (319) Lee, C. L., et al. 1964. Effect of papain on antigenicity of mouse erythrocytes. In: 48th Annual meeting of the Federation of American Societies for Experimental Biology, Chicago, 1964. Fed. Proc. 23 (2Pt.1): 389
- (320) Leipert, T., and I. Häfner. 1930. Über die Abspaltung von Tyrosin und Tryptophan aus Casein durch mit Blausäure aktiviertes Papain. Biochem. Ztschr. 229: 427-432
- (321) Leont'eva -Tyukina, A. A. 1967. Elastase activity of papain. Profl. Travmatizma, Lecheniya Travm Zabol. Oporno-Dvigat. App. 337-341
- (322) Leslie, R. G., et al. 1971. The products from papain and pepsin hydrolyses of guinea-pig immunoglobulins gamma 1G and gamma 2G. Biochem. J. 121: 829-837

- (323) Lillov, L., et al. 1969. Influence of some plant proteases on liver tissue. VI. Influence of ficin, papain and bromelain on the cytochrome oxidase activity of liver mitochondria. *Research Workers* 15: 389-396
- (324) Lineweaver, H., and S. R. Hoover. 1941. A comparison of the action of crystalline papain on native and urea-denatured proteins. *J. Biol. Chem.* 137: 325-335
- (325) Loo, Y. H., et al. 1960. Erythropoietic activity of proteolytic enzymes. *Proc. Soc. Exptl. Biol. Med.* 104: 39-40
- (326) Lopez, A., et al. 1955. Enzymic inhibition of gelation in frozen egg yolk. *J. Milk and Food Technol.* 18: 77-80
- (327) Lorand, L., and K. Konishi. 1962. Protein-polymerizing activity of papain. *Biochem. Biophys. Res. Commun.* 7: 457-460
- (328) Lowry, R. J., et al. 1956. Lysozyme-induced sensitivity of *Neurospora* ascospores to furfural. *Mycologia* 48(2): 241-252
- (329) Lunt, W. R. F. 1934. Verbessern von Nahrungsmitteln. *Brit. P.* 409790
- (330) Luscombe, M., and C. F. Phelps. 1967. Action of degradative enzymes on the light fraction of bovine septa protein polysaccharide. *Biochem. J.* 103: 103-109
- (331) Maher, F. T., and E. H. Wirth. 1946. The assay of papain with suggestions for the modification of the tentatively accepted assay. *Bull. Natl. Formulary Comm.* 14: 56-70
- (332) Maier, C. L. Jr. 1968. Preparative thin-layer gel filtration. *J. Chromatogr.* 32(3) 577-579

- (333) Mandy, W. J. 1967. A new serum factor in normal rabbits. III. Specificity for antigenic determinants uncovered by papain or pepsin digestion. J. Immunol. 99(4): 815-824

- (334) Mangiarotti, G., et al. 1963. Alteration of enzymatic activities of rat liver diphosphatase by treatment with papain or urea. Biochem. Biophys Res. Commun. 12: 305-309

- (335) Marks, C., et al. 1964. The effects of papain on the skull of the rat. J. Dent. Res. 43(5 Pt.2): 758-759

- (336) Markwardt, F., et al. 1965. Mechanism of the amine-releasing effect of proteolytic enzymes in thrombocytes. Arch. Exptl. Pathol. Pharmacol. 251(3): 255-264

- (337) Mars. P. H. 1954. Assay of papain preparations. Pharm. Weekblad. 89: 97-98

- (338) Maschmann, E., and E. Helmert. 1934. Activation by vitamin C (ascorbic acid) or by vitamin C iron. Ztschr. f. physiol. Chem. 224: 56-60

- (339) Maschmann, E., and E. Helmert. 1934. Activation of papain by vitamin C-iron and restriction of its activity by vitamin C. Ztschr. f. physiol. Chem. 223: 127-135

- (340) Maschmann, E., and E. Helmert. 1935. Activation and inhibition on intracellular proteinase: further studies. Biochem. Ztschr. 280: 184-203

- (341) Mateyko, G. M. 1967. Developmental modifications in *Arbacia punctulata* by various metabolic substances. Biol. Bull. 133(1): 184-228

- (342) Matsumoto, H., and A. Kanatani. 1953. Breadmaking. VIII. Measurement of the pressure extrusion viscosity of dough. J. Fermentation Technol. 31: 485-487

- (343) Mazza, F. P., and C. Migliardi. 1942. The industrial application of papain digestion in the extraction of tunny-liver oil. *Ricerca Sci. e Ricostruzione* 13: 267-270
- * (344) Mazzei, D., et al. 1966. Mitogenic action of papain. *Lancet* 2: 802-803
- (345) McDonald, K. G., and S. K. Sim. 1957. Activation of histamine release from rabbit blood by crystalline papain. *Can. Pharm. J.* 90: 231-234
- (346) McElligott, T. F., and J. L. Potter. 1960. Increased fixation of sulfur-35 by cartilage in vitro following depletion of the matrix by intravenous papain. *J. Exp. Med.* 112: 743-750
- (347) McFarlane, W. D., et al. 1963. Retention of clarity, color, and flavor in beer through control of the tannin content. *Am. Soc. Brewing Chemists.* 1963, 98-103
- (348) McGinnis, J., and V. H. Menzies. 1946. Effect of in vitro enzymatic digestion of raw soybean flakes on chick growth. *Poultry Sci.* 25(5): 538-539
- * (349) McGraw, J. Y. 1964. Interrelation of serum cholesterol level, aorta cholesterol content, and the degree of atherosclerosis in experimental cholesterol atherosclerosis. *Laval Med.* 35(8): 909-917
- (350) McVaugh, R. 1943. To make tough meat tender. *Agriculture in the Americas.* 3(7): 134-136
- (351) Medgyesi, G. A., et al. 1970. Papain sensitivity of 7S mouse myeloma proteins. *Progr. Immunobiol Stan.* 4: 71-76
- * * (351a) The Merck Index. 1968. 8th. Ed. Merck and Co. Rahway, New Jersey. p.781

- (352) Merkow, L., and J. J. Lalich. 1961. Skeletal changes in suckling rats induced by prolonged papain administration. J. Bone and Joint Surg. 43A(5): 679-686
- (353) Mihaescu, C., and M. Seligmann. 1968. Papain digestion fragments of human IgM globulins. J. Exp. Med. 127(3): 431-453
- (354) Miller, B. S. 1949. Report on (the determination of) proteolytic activity of flour and other proteolytic enzyme-containing materials. J. Assoc. Offic. Agr. Chemists. 32: 261-267
- (355) Minker, E., and M. Koltai. 1967. The ganglionic effect of trypsin. Acta Physiol. Acad. Sci. Hung. 31(2): 167-176
- (356) Mokrasch, L. C., and R. W. McGilvery. 1956. Purification and properties of fructose-1, 6-diphosphatase. J. Biol. Chem. 221: 909-917
- (357) Moll, M., et al. 1971. Some assay methods for proteolytic enzymes. BIOS (3) 3-4 & 7-11
- * (358) Monkhouse, F. C. 1955. An investigation of the coagulation defect brought about in rabbits and dogs by intravenous injection of papain. Canadian J. Biochem. and Physiol. 33(2): 112-121
- (359) Moreau, J. R., and E. E. Jankus. 1963. An essay for measuring papain in meat tissue. Food Technol. 17(8): 93-95
- (360) Morgan, W. T. J. 1959. Immunochemical aspects of the products of the human blood-group genes. Ciba Foundation Symposium, Biochemistry of Human Genetics 1959: 194-216
- (361) Morgan, W. T. J., and A. Pusztai. 1961. Studies in immunochemistry. 21. The products of the degradation of blood-group-specific substances by ficin and papain. Biochem. J. 81(3): 648-658

- (362) Morris, M., and C. V. Seastone. 1955. The relation of M protein and resistance to phagocytosis in the beta hemolytic streptococci. J. Bacteriol. 69: 195-203
- (363) Morton, J. A. 1962. The action of blood-group antibodies on red cells treated with proteolytic enzymes. Brit. J. Haematol. 8: 134-148
- (364) Murata, Y. 1971. Experimental studies on the influence of papain on the intervertebral disk. J. Jap. Orthop Assoc. 45: 257-266
- (365) Murray, D. G. 1964. Experimentally induced arthritis using intra-articular papain. Arth. and Rheum. 7(3): 211-219
- (366) Murray, M., et al. 1965. The enzymic induction of atherosclerosis. Arch. Pathol. 79(2): 144-162
- (367) Murray, S. 1971. Early "enzyme" Rh antibodies and second pregnancies. Vox Sang 21: 217-221
- (367a) NAS/NRC. 1965. Chemicals used in Food Processing. Publication 1274. p.267
- (367b) NAS/NRC, 1972. Comprehensive Gras Survey. Table II Part A, p.7; Table 2, p.24; Table 13, Part A p.118
- (368) Nakashima, R. 1927. Über ein Tetrapeptid aus Gliadin J. Biochemistry 7: 441-446
- (369) Nanikawa, R., et al. 1963. Studies on experimental cardiac necrosis produced by means of intravenous administration of proteolytic enzymes. The changes of the activities of some serum enzymes and the histological findings of several organs in papain injected rabbits. Jap. J. Leg. Med. 17: 304-313

- (370) Naumovski, A., and B. Nikodijevic. 1960. Intra-peritoneal administration of papain. *Acta Pharm. Jugoslav.* 10: 79-84
- (371) Netherlands Patent. 1966. 6 515 289. Anonymous. Foods.
- (372) Nihei, T., and C. M. Kay. 1968. Isolation and properties of an enzymically active fragment from papain-digested myosin. *Biochem. Biophys. Acta* 160(1): 46-52
- (373) Nogami, H., et al. 1960. Acid-stable digestive enzyme. IV. Comparison of digestibility with various proteases. *Yakugaku Zasshi.* 80: 1788-1794
- (374) Nordberg, B. K., and W. Thorsell. 1955. The effects of certain enzyme systems on the capsule of *Bacillus anthracis*. *J. Bacteriol.* 69: 367-371
- (375) Northrop, J. H. 1947. Action of papain and ficin on tadpoles and *Arbacia* eggs. *J. Gen. Physiol.* 30(5): 375-376
- (375a) OPD Chemical Buyers' Directory. 1970-1971. Schnell Publishing Co., Inc. New York. p. 1020
- (376) Odintsov, A. I. 1957. Enzymes used in cooking. *Voprosy Pitaniya* 16(2): 73-76
- (377) Oken, D. E., and R. J. Boucek. 1957. Enzymatic proteolysis of rat sponge biopsy collagen tissue. *Proc. Soc. Exptl. Biol. and Med.* 96(2): 367-369
- (378) Olitzki, A. L., and D. Sulitzeanu. 1958. Studies on the antigenic structure of *Brucella suis* with the aid of the agar gel precipitation technique. I. The resistance of antigens to physical, chemical and enzymatic treatments. *Brit. J. Exptl. Path.* 39(3): 219-227

- (379) Onoue, K., et al. 1967. Papain fragmentation of the subunits of human macroglobulin. J. Immun. 98: 303-313
- (380) Otaka, F., and F. Hongo. 1965. Digestion of beef by various proteolytic enzymes. Nippon Shokuhin Kogyo Gakkaishi 12(3): 101-105
- (381) Parke, J. A. C., and P. J. Avis. 1964. The effect of digestion with papain and pepsin upon the antioxic of rabbit antibody. Immunology 7(3): 248-260
- (382) Partmann, W. 1955. Artificial ripening of meat with papain preparations. Z. Lebensm-Untersuch. u.-Forsch. 102: 321-329
- (383) Penau, H., and R. Audic. 1934. Matières passives normales pour le dosage des ferments solubles pharmaceutiques. J. Pharm. et Chim. 19(7): 329-345
- (384) Penau, H., and R. Audic. 1937. Assay of soluble normalized enzymes and substrates. J. Pharm. Chim. 25: 107-110
- (385) Penny, I. F. 1960. Up-grading of low - grade meat. Chem. and Ind. 288-289
- (386) Perona, G., et al. 1965. Loss of acetylcholinesterase in human erythrocytes treated with trypsin, papain or bromelin; its relation with susceptibility to acid lysis in vitro. Brit. J. Haematol. 11(2): 171-176
- (387) Petermann, M. L. 1942. On beef serum pseudoglobulin and on diphtheria anti-toxin. J. Biol. Chem. 144: 607-616
- (388) Petermann, M. L. 1946. Splitting of human gamma globulin antibodies by papain and bromelin. J. Am. Chem. Soc. 68: 106-113

- (389) Petrovskii, Y. A. 1936. The mechanism of the action of a series of medicinal agents on the biliary tract. *J. Physiol. (U.S.S.R.)* 21: 293-310
- (390) Pettijohn, O. G. 1940. A study of the oxidation-reduction potentials of Ohio soft red winter wheat flours and doughs. *Ohio State Univ. Abst. of Doctor's Dissert.* 33: 199-205
- (391) Piukovich, I., et al. 1968. Serum glycoprotein levels in papain-treated hepatectomized rats. *Kiserl. Orvostud.* 20(4): 436-441
- (392) Podrouzek, V. 1942. Methods for proteolytic systems. *Casopis Ceskoslov Lekarnictva.* 21: 171-177
- (393) Pogell, B. M., and R. W. McGilvery. 1952. The proteolytic activation of fructose-1, 6-diphosphatase. *J. Biol. Chem.* 197: 293-302
- (394) Poljak, R. J., et al. 1966. Papain fragmentation of rabbit gamma-globulin: a crystallographic study. *J. Molec Biol.* 17: 546-547
- (395) Pontremdi, B., et al. 1965. Fructose diphosphatase from rabbit liver. IV. Sulfhydryl groups and their relations to the catalytic activity. *Arch. Biochem. Biophys.* 112(1): 7-15
- (396) Potter, J. L., et al. 1960. The effects of papain on cartilage in vivo: Factors influencing the distribution of papain protease following intravenous injection. In: *Connective tissue and diseases of connective tissue.* Ann. N. Y. Acad. Sci. 86(4): 929-942
- (397) Potter, J. L. 1961. Papain and flop-eared rabbits. *Arthritis Rheumatism* 4: 389-399
- (398) Potter, J. L., et al. 1962. Impairment of enzyme-binding capacity of serum in rheumatoid disease. *Proc. Roy. Soc. Med.* 55: 111-113

- (399) Pozsar, H. K., et al. 1969. Investigations of milk clotting enzymes from the point of view of cheese production. III. Selective inhibition of the proteolytic activity of plant proteases. *Élelmiszertudomány*. 3(2): 55-70
- (400) Pozerski, E. 1920. Wirkung des Papains auf das *Bacterium coli*. *C. r. Soc. de Biologie*. 83: 751-753
- (401) Prager, M. D., et al. 1962. Mechanism of the immunohematologic effect of papain and related enzymes. *J. Immunol.* 89: 834-840
- (402) Pratolongo, U. 1938. Biochemische Probleme und Untersuchungen in der Technik der Brotbereitung. *Chim. e Ind. (Milano)* 20: 129-132
- (403) Pressman, R. 1957. Coatings for meat-wrapping sheets. *U. S. Patent* 2 811 454
- (404) Prioreschi, P. 1964. Role of sodium and potassium in the cardiopathy produced by papain. *Path. Biol. (Paris)* 12: 206-208
- (405) Purr, A., and M. Russell. 1934. Die Aktivierungsfähigkeit des Papains, angewendet auf eine Bestimmungsmethode physiologisch aktiver Stoffe in Blut. *Hoppe-Seyler's Z. Physiol. Chem.* 228: 198-206
- (406) Purr, A., and M. Lubieniecki-von Schelhorn. 1970. Studies of the mechanism of the effect of hexamethylenetetramine and glyoxylic acid on the proteolysis and decarboxylation of amino acids in sour fish preparations. *Deutsche Lebensmittel-Rundschau*. 66(4): 111-115
- * * (407) Pushpakom, R., et al. 1970. Experimental papain-induced emphysema in dogs. *Am. Rev. Resp. Dis.* 102: 778-789
- (408) Pusztai, A., and W. T. J. Morgan. 1958. Degradation of blood group-specific mucopolysaccharides by ficin and papain. *Nature* 182(4636): 648-650

- (409) Pusztai, A., and W. T. J. Morgan. 1961. Immunochemistry. XX. The action of papain and ficin on blood-group-specific substances. *Biochem. J.* 81: 639-647
- (410) Ramon, G. 1943. Effects, ferment, anaferment. antiferment. *Compt. rend Acad. d. sc.* 217: 562
- (411) Ramon, G. 1945. Ferment, anaferment, antiferment; introduction to comparative immunologic study with toxins, anatoxins antitoxins, viruses, ana-viruses. *Bull. Acad. de med.* 129: 8-13
- (412) Ramon, G. 1945. Effects, ferment, anaferment, antiferment. *Rev. d'immunol.* 9: 134-138
- (413) Randelli, G., et al. 1962. Changes in the growth zone caused by papain. *Arch. Ortop.* 75: 1405-1413
- (414) Rao, V. K. M., et al. 1952. Enzyme inhibition studies in relation to drug action. II. Action of certain antibiotics on papain. *J. Sci. and Indust. Res. (India)* 11B(7): 299-302
- (415) Ray, N. N. 1938. Zerstörung verschiedener Toxine durch proteolytische Enzyme. *Sci. and Cult.* 3: 496
- (416) Read, J. W., and L. W. Haas. 1934. Die Backfähigkeit der Mehle als Einwirkung verschiedener enzymatischer Eingriffe. *Cereal Chem.* 11: 280-298
- (417) Read, R. O. 1970. Keeping up with all the recent changes in Food and Drug Regulations. *Canadian Food Industries* 41(3): 39-41
- (418) Reigier, L. W., and A. L. Tappel. 1956. Freeze-dried meat. III. Non-oxidative deterioration of freeze-dried beef. *Food Res.* 21(6): 630-639

- (419) Rich, A. R., and G. L. Duff. 1937. Die Erzeugung von hyaliner Arteriosklerose und Arterioneckose mittels proteolytischer Enzyme. Bull. Johns Hopkins Hosp. 61: 63-70
- (420) Remy, R., and P. Lallouette. 1969. Action of normal serum and antipapain serum of the rabbit against acute inflammatory reactions induced in the mouse by papain. C. R. Acad. Sci. Ser. D. 269(22): 2280-2282
- (421) Rigaud, F., et al. 1949. A new aspect of the properties of papain: bacteriostasis. Compt. Rend. Soc. Biol. 143(5/6): 399-400
- (422) Rigaud, F., et al. 1950. A new aspect of the properties of papain: bacteriostasis. J. Med. Bordeaux. 127(1): 42-43
- (423) Ringer, W. E., and B. W. Grutterink. 1926. Influence of reaction on papain digestion of protein. Ztschr. Physiol. Chem. 156: 275-324
- (424) Ringer, W. E., and B. W. Grutterink. 1927. Einfluß der Reaktion auf die eiweißverdauende Kraft des Papains. II. Mitt. Ztschr. Physiol. Chem. 164: 112-142
- (425) Ringer, W. E. 1930. Papain and its effects. Arch. di farmacol. sper. 48: 99-116
- (426) Robert, L., et al. 1961. Serum haptoglobin in experimental connective tissue disorders. Proc. Soc. Exptl. Biol. Med. 107: 499-502
- (427) Robert, L., and B. Robert. 1963. Enzymic mechanisms of normal and pathological catabolism of connective tissue. Exposés Ann. Biochim. Med. 24: 269-290
- (428) Robinson, F. A. 1945. "F" food. Food Manufacture 20(8): 294-298

- (429) Roche e Silva, M., and S.O. Andrade. 1943. Histamine and proteolytic enzymes--liberation of histamine by papain. J. Biol. Chem. 149: 9-17
- (430) Roche e Silva, M., et al. 1946. Chemical aspects on the liberation of histamine from tissues. Selecta Chim. No. 4: 65-84
- (431) Rogers, W., et al. 1965. Effects of enzyme addition on palatability, tenderness, and rancidity development in quick-aged country-style hams. Food Technol. 19(10): 1590-1593
- (432) Rogers, R. W. 1970. The effect of pancreatic lipase and papain on palatability, tenderness and rancidity development of quick-aged country-style hams. Dissertation Abstracts International. Section B. 31(4): 1629
- (433) Rondoni, P. 1950. Action of hydrocarbons and other substances, carcinogenic or not, on SH enzyme activity. Ricerca Sci. 20: 671-672
- (434) Rosenheim, A. H. 1937. The action of enzymes on antibodies. Biochem. J. 31: 54-71
- (435) Roth, G. D., and R. J. Bruckner. 1960. The action of enzymes on ethylene oxide-treated periodontal tissue of rats. J. Dent. Res. 39(1): 76-83
- (436) Roth, J. S., et al. 1960. Function of intracellular ribonucleases. II. Interaction of ribonucleoprotein and enzymes. J. Biophys. Biochem. Cytol. 7: 443-453
- (437) Rotsch, A. 1966. Application of enzymes in cracker production. Brot. Gebaeck. 20(11): 213-217
- (438) Rowe, D. S. 1962. The separation and antigenic characteristics of some fractions of papain digests of human serum gamma globulins with the gamma globulins of some other mammalian sera. Immunology. 5(4): 533-548

- (439) Rubinstein, H. M. 1957. Coagulant action of proteolytic enzymes. *Nature* 180: 1202-1203
- (440) Ruffolo, P. R. 1964. Correlated light and electron microscopic study of myocardial necrosis induced in the rat by intravenous papain. In: *Scientific Proceedings of the American Association of Pathologists & Bacteriologists, Chicago, 1964*. Amer. J. Pathol. 44(6): 26a-27a
- (441) Ruffolo, P. R. 1964. The Pathogenesis of necrosis, I. A correlated light and electron microscopic observations of the myocardial necrosis induced by the intravenous injection of papain. *Amer. J. Pathol.* 45(5): 741-756
- (442) Rupp, E., and C. H. Bailey. 1937. Die Wirkung von Proteasezusätzen auf die plastischen Eigenschaften der Teige. *Cereal Chem.* 14: 496-510
- (443) Sachdev, J. C. 1954. Antifatty liver action of proteolytic enzymes of animal and plant origin in pancreatic duct-ligated rats. *Gastroenterology*. 27: 353-357
- (444) Sanchez-Avalos, J., and S. P. Miller. 1968. Degradation of fibrinogen by proteolytic enzymes. I. Characterization and isolation of products and their relation to fibrinolytic states. *Thromb. Diath. Haemorrh.* 19(3-4) 499-515
- (445) Sandstedt, R. M. 1945. The relation of proteolysis to the characteristics of oxidation and reduction in doughs. II. Some reasons for conflicting interpretations of data. *Baker's Digest*. 19(5): 125-126
- (446) Sato, S. 1929. Physico-chemical investigation on the casein-splitting action of papain with special reference to the fundamental properties of its action as well as to the mechanism of the acceleration of HCN solution upon its action. *J. Fac. Agric. Hokkaido Imp. Univ.* 24(4): 101-151

- (447) Sato, T., et al. 1962. Utilization of protease in baking. I. Proteolytic activities of various kinds of protease and their effects on baking. Nippon Shokuhin Kogyo Gakkaishi. 9(8): 326-331
- (448) Saunders, A. M., et al. 1967. Electron microscopy of chondromucoprotein and the products of its digestion with hyaluronidase and papain. Nature (London) 214: 194-195
- (449) Savage, D. J., and C. C. Thompson. 1971. The isoelectric focussing of beer proteins in polyacrylamide gel. J. of the Institute of Brewing. 77(4): 371-375
- (450) Schales, O., et al. 1948. Inhibition of enzymatic proteolysis. I. Observations with carbonyl group reagents; effect of hydrazine on peptic hydrolysis. Arch. Biochem. 19(1): 119-132
- (451) Schally, A. V., et al. 1971. Further studies on the enzymatic and chemical inactivation of hypothalamic FSH-RH. Neuroendocrinology 8: 347-358
- (452) Scharpenseel, H. W., and F. C. Anastacio. 1955. Farm-raised poultry-feeding supplements. Araneta J. Agr. 2(4): 16-41
- (453) Scharpenseel, H. W. 1956. Farm grown fodder supplements for hens and their significance for growth promotion and saving of animal protein. Landwirtsch. Forsch. 8: 219-226
- (454) Schecter, I., et al. 1967. On the size of the active site in proteases. I. Papain. Biochem. Biophys. Res. Commun. 27: 157-162
- (456) Schiff, W., and E. Seidlitz. 1954. A hemagglutination test for tuberculosis (Middlebrook-Dubos) with papanized blood cells. Zeitschr. Immunitätsforsch. 111(2): 69-89
- (457) Schmidt, H. 1958. The effects of fluorine-containing and surface-active substances on the enzymic degradation of dentine. ORCA, Communs. Congr. organisme européen coord. recherches Fluor prophylaxie carie dentaire 5th, Brussels, Belg. 1958: 240-251

- (458) Schneider, M., et al. 1965. Assay of papain using ^{131}I -labeled albumin. Proc. Soc. Exptl. Biol. Med. 120(1): 130-134
- (459) Schneider, M., and G. Goldstein. 1967. Digestion of articular cartilage by papain in vitro. Proc. Soc. Exp. Biol. Med. 124(3): 991-994
- (460) Schwartz, B. D., et al. 1971. Regeneration of transplantation antigens on mouse cells. Transplant Proc. 3: 180-182
- (461) Sciban, R., et al. 1969. Role of activators in proteolytic treatment of beers. Tageszeitung für Brauerei. 66(85/86): 554-556
- (462) Scriban, R., and M. Stienne. 1970. Detection and quantitative determination of papain in beer. J. of the Institute of Brewing. 76(3): 243-244
- (463) Scriban, R., et al. 1970. Role of activators in proteolytic treatment of beers. Proceedings. European Brewing Convention 12: 397-417
- (464) Seaman, G. V. F., and G. Uhlenbruck. 1962. Action of proteolytic enzymes on the red cells of some animal species. Biochim. Biophys. Acta 64: 570-572
- (465) Seaman, G. V. F., and G. Uhlenbruck. 1962. Electrophoretic mobility of erythrocytes after treatment with various enzymes and antiserum. Klin. Wochschr. 40: 699-701
- (466) Selye, H., and E. Bajusz. 1958. Prevention by MgCl_2 and KCl of the myocardial necroses normally produced by papain. Cardiologia 33(4): 305-312
- (467) Seyle, H., and E. Bajusz. 1959. Sensitization by K or Mg deficiency for the production of myocardial necroses by papain. Z. Vitamin-, Hormon- u. Fermentforsch. 10: 39-45

- (468) Selye, H., and E. Bajusz. 1959. The age factor in the production of various experimental cardiopathies. J. Gerontol. 14(2): 164-168
- (469) Selye, H., and P. Rohan. 1966. Humoral conditioning for necrosis. Naunyn-Schmiedbergs Arch. Pharmakol. Exp. Pathol. 255(2): 142-150
- (470) Sen, D. P., et al. 1962. Fish Hydrolysates, I. Rate of hydrolysis of fish flesh with papain. Food Technol. 16(5): 138-141
- (471) Shapira, E., and R. Arnon. 1967. The mechanism of inhibition of papain by its specific antibodies. Biochemistry 6(12): 3951-3956
- (472) Sheldon, H., and R. A. Robinson. 1960. Studies on cartilage. II. Electron microscope observations on rabbit ear cartilage following the administration of papain. J. Biophys. Biochem. Cytol. 8: 151-163
- (473) Shibata, Y. 1966. Tryptic digestion of human γ -globulin. Seikagaku 38: 874-882
- (474) Shinke, R., and N. Mugibayashi. 1969. Studies on barley malt amylases. XIV. Mechanism of activation of zymogen β -amylase by proteolytic enzymes and SH compounds. J. of the Agricultural Chemical Society of Japan. 43(5) 278-285
- (475) Simms, H. S., and N. P. Stillman. 1937. Stoffe, welche reifes Gewebe in vitro beeinflussen. I. Die stimulierende Wirkung von Trypsin auf frisches reifes Gewebe. J. gen. Physiol. 20: 603-627
- (476) Singh, S. I., and I. Singh. 1955. Active relaxation of unstriated muscle produced by proteolytic enzymes and the mode of action of these enzymes. Proc. Indian Acad. Sci. Sect. B. 42(3): 85-89

- (477) Sinha, R. C., and K. Bahadur. 1958. Comparative study of the proteolytic activities of papain and trypsin against casein, egg albumin, and meat extract. *Enzymologia* 19: 319-322
- (478) Skelton, G. S. 1971. The relation between "milk clotting" and proteolytic activity in papain studies. *Enzymologia* 40(3): 170-172
- (479) Sluyterman, L. A. E. 1966. The relation between enzyme, substrate, and product. *Philips Tech. Rev.* 27(6): 160-168
- (480) Sluyterman, L. A. AE. 1966. Substrate binding by non-activated papain. *Biochim. Biophys. Acta.* 113(3): 557-586
- (481) Sluyterman, L. A. 1967. The activation reaction of papain. *Biochim. Biophys. Acta.* 139: 430-438
- (482) Sluyterman, L. E. A., and B. G. Wolthers. 1969. A tentative mechanism of papain action. *Proc. Kon. Ned. Akad. Wetensch Ser. B. Phys. Sci.* 72(1): 14-16
- (483) Smallings, J. B., et al. 1971. Effect of antemortem injection of papain on the tenderness and quality of dry-cured hams. *J. of Animal Science.* 32(6): 1107-1112
- (483a) Smith, E. L. and J. R. Kimmel, 1960. Papain (with a section on ficin) In: P.D. Boyer, et al., eds. *Enzymes.* Academic Press. Vol. 4: 133-173
- (484) Smith, E. L., et al. 1954. Crystalline papain. II. Physical studies; the mercury complex. *J. Biol. Chem.* 207(2): 533-549
- (485) Smith, E. L., et al. 1954. Crystalline papain III. Amino acid composition. *J. Biol. Chem.* 207(2): 551-561

- (485a) Smith, E. L. 1958. Active sites of papain and covalent "High-Energy" bonds of protein. J. Biol. Chem. 233: 1392
- (486) Smorodintsev, I. A., and V. P. Zhigalov. 1941. Influence of papain and pepsin on proteins of meat and gluten. Compt. rend. acad. sci. U.S.S.R. 33: 70-72
- (487) Solov'ev, V. I., and L. G. Gashimova. 1970. Effect of papain on the organoleptic and physicochemical properties of buffalo meat. Prikl. Biokhim. Mikrobiol. 6(2): 190-195
- (488) Sonenberg, M., and J. M. Dellacha. 1967. Anabolic effects in man of papain digests of bovine growth hormone. J. Clin. Endocrinol. Metab. 27(7): 1035-1040
- (489) Sosebee, M. E., et al. 1964. The effects of enzyme addition on the quality of freeze-dehydrated chicken meat. Food Technol. 18(4): 149-152
- (490) Spanish Patent. 1954. 214226. Poch, Maria Teresa de la Puente. Tenderizing of meat.
- (491) Spanish Patent. 1958. 239494. Halor Sociedad Limitada. Meat tenderizer
- (492) Spicer, S. S., and J. H. Bryant. 1957. Cartilage changes in papain-treated rabbits. Amer. J. Pathol. 33(6): 1237-1245
- (493) Spicer, S. S., and J. H. Bryant. 1958. Systemic effects in rabbits receiving injections of papain and chondroitin sulfate. Am. J. Pathol. 34: 61-75
- (494) Spielmann, W. 1954. On influencing the conglutinated anti-Rh sera with different chemicals. Part V. Zeitschr. Immunitätsforsch. 111(5): 470-484

- (505) Tachibana, T. 1942. Influence of proteinases upon various toxins. I. Diphtheria and tetanus bacterial toxins. Japan J. Microbiol. and Path. 36: 220-236
- (506) Tachibana, T. 1942. Influence of proteinases upon the endotoxins of cholera vibrio and typhoid bacillus. Japan. J. Microbiol. Pathol. 36: 493-503
- (507) Takahashi, J. 1970. Methods of quick ripening of slated viscera utilizing the viscera of swine. J. of Food Science Technology. (Nihon Shokuhin Kogyo Gakkai-shi) 17(9): 412-415
- * * (508) Takaoka, T. 1955. Some toxicological and pharmacological properties of an enzymic anthelmintic. Vermizym. Folia Pharmacol. Japan. 51: 27-33
- (509) Takeuchi, J. 1964. Effect of papain on the growth of chicken sarcoma in vivo. Gann 55(3): 259-261
- (510) Takeuchi, T. 1960. Induction of mitosis in smooth muscle fibers of coronary artery of an adult rabbit by means of tissue protease. Mie Med. J. 10: 9-16
- (511) Tappel, A. L., et al. 1956. Meat tenderization. II. Factors affecting the tenderization of beef by papain. Food Res. 2(3): 375-383
- (512) Tassoni, G. 1970. Further studies in vitro of the fluidifying action of enzymes, reducing substances and their association. Valsaiva. 46: 281-285
- (513) Tauber, H. 1935. Aktivatoren und Hemmungskörper von Enzymen. Ergebn. Enzymforsch. 4: 42-67
- (514) Tazawa, Y. 1943. Cleavage of protamines by proteinases. Acta Phytochim. (Japan) 13: 139-153

- (495) Springer, R., and K. E. Quentin. 1955. Studies on protein sulfur. V. The behavior of protein sulfur by enzymic degradation of albumin. Arch. Pharm. 288: 448-452
- (496) Ssadikow, W. 1926. "Über die Eigenschaften von Haut-und Sehnenkollagen und über ihr Verhalten gegenüber Fermenten Collegium. 1926, 512-518
- (497) Steiner, R. F., and K. Laki. 1951. Light-scattering studies on the clotting of fibrinogen. Arch. Biochem. Biophys. 34: 24-37
- (498) Stern, P., et al. 1968. Pharmacological analysis of a mucopolysaccharide polysulfuric acid ester. Z. Rheumaforsch, 27(7-8): 254-260
- (499) Stern, R. 1948. Comparative study of the effects of cysteine hydrochloride and papain on unsalted and salted doughs. Cereal Chem. 25: 220-228
- (500) Stevens, L. E. 1968. A reassessment of papain in preventing peritoneal adhesions. Amer. J. Surg. 115(4): 535-539
- (501) Stupenengo, R. H., et al. 1964. Antitussive drugs. Semana Med. 125(10): 391-395
- (502) Stylos, W. A., and N. R. Rose. 1968. Rabbit thyroid auto-antibody produced by intravenous injections of papain-treated rabbit thyroid extract. Proc. Soc. Exp. Biol. Med. 127(3): 784-786
- (503) Suzuki, H. K. 1966. Localized hyperchondroplasia induced by crude papain administered to mice. Nature (London) 209: 1111-1113
- (504) Svedberg, T., and I. B. Erikson. 1933. Abbau des Ovalbuminmolekuls durch Papain. Biochem. Z. 258: 1-3

- (515) Telegdy-Kovats, L., and M. Kelemen. 1965. Enzyme treatment of beef. *Nahrung* 9(3): 299-306
- (516) Thomas, L. 1956. Reversible collapse of rabbit ears after intravenous papain, and prevention of recovery by cortisone. *J. Exptl. Med.* 104: 245-251
- (517) Thomas, L., et al. 1960. Comparison of the effects of papain in vitamin A on cartilage. I. The effects in rabbits. *J. Exp. Med.* 111: 705-718
- (518) Thomas, L. 1964. The effects of papain, vitamin A, and cortisone on cartilage matrix in vivo. *Bio-phys. J.* 4(1): 207-213
- (519) Thompson, E. O. P. 1954. Crystalline papain IV. Free amino groups and nitrogen terminal sequence. *J. Biol. Chem.* 207(2): 563-574
- (520) Thompson, R. F. 1941. Report on (the assay of) enzymes (papain). *J. Assoc. Official Agr. Chem.* 24: 540
- * (521) Thompson, S. W. II 1957. Toxicity of various enzymes when inoculated intracerebrally into mice. *Cornell Vet.* 47(2): 326-334
- (522) Thorpe, N. O., et al. 1966. Studies on papain produced subunits of human gamma-G-globulins I. Physicochemical and immunochemical properties of gamma-G-globulin Fc-fragments. *Immunochemistry.* 3: 317-327
- (523) Tobias, J. M. 1955. Effects of phospholipases, collagenase and chymotrypsin on impulse conduction and resting potential in the lobster axon with parallel experiments on frog muscle. *J. Cell. and Comp. Physiol.* 46(2): 183-208
- (524) Tomcsik, J., and J. B. Baumann-Grace 1957. Action of trypsin on bacterial cell structures. *Schweiz. Z. allgem. Pathol. u. Bakteriologie.* 20: 129-144

- (525) Tomcsik, J., and J. B. Baumann-Grace. 1960. Action of proteolytic enzymes on the mononucleosis antigen in sheep and beef erythrocytes. Pathol. et Microbiol. 23: 172-183
- (526) Tsaltas, T. T., and K. A. Greenawald. 1964. Papain-induced changes in rabbit cartilage. II. Chemical changes in the cartilage matrix of "adolescent" rabbits. Clin. Orthop. Related Res. 38: 170-178
- (527) Tsaltas, T. T., and C. E. Lutton. 1965. Effect of intravenous papain on the plasma lipids and fatty acids of adult rabbits. J. Atherosclerosis Res. 5(6): 533-539
- (528) Tsaltas, T. T., and K. A. Greenawald. 1966. Comparison of changes in the chemical composition of rabbit cartilage matrix with age and following intravenous papain. Nature 210(5042): 1237-1239
- (529) Tsou, C. L. 1951. Cytochrome c modified by digestion with proteolytic enzymes. I. Digestion. Biochem. J. 49: 362-367
- (530) Tsugo, T., and K. Yamauchi. 1955. Milk-coagulating enzymes. IV. Inhibition of milk-coagulating enzymes by blood serum. Nippon Nogei-kagaku Kaishi. 29: 457-461
- (531) Tsuk, A. A., and G. Oster. 1961. Determination of enzyme activity by a linear measurement. Nature 190: 721-722
- (532) Underrain, A. 1933. ¹¹Über die Wirkung des Papayotins auf die Milch. Forschg. 15: 433-447 Milchwirtschaftl.
- (533) Ungar-Waron, H., et al. 1967. Action of papain on normal rabbit immuno-globulin M. (Carcia-papaya-D.) Biochim Biophys. Acta. 140(3): 542-545
- * (533a) The U.S. Dispensary 1973. 27th. Ed. J. B. Lippincott Co. Philadelphia, Pa. p. 829

- (547) U.S. Patent. 1971. 3 561 975. Luebering, J. F.,
and E. R. Hair. Pie dough improvement.
- (548) Unwin, E. E. 1937. Papain. Its assay. Chemist
and Druggist. 127: 399-400
- (549) Van Eyke, H. G. 1966. Fragmentation of human globulin
with papain. Biochim. Biophys. Acta 127(1):
241-243
- (550) Van Laer., H. 1919. Reaktionen zwischen Enzymen.
Bull. Soc. Chim. Belgique 28: 340-346
- (551) Velluz, L. 1938. Inactivation de quelques toxines
par la papaine. Compt. Rend. Soc. Biol. 128(16):
132-133
- (552) Velluz, L. 1938. Effects of inactivation of strychnine
sulfate. Compt. rend. Soc. de Biol. 128: 291-293
- (553) Visco, S. 1926. Über die physikalisch-chemischen
Veränderungen der Eiweißhydrosole durch einige
pflanzliche Milchsäfte. Arch. di Scienze biol.
9: 41-73
- (554) Voegtlin, C., et al. 1933. The influence of the
oxygen tension on the reversal of proteolysis
(protein synthesis) in certain malignant tumors
and normal tissues. J. Pharmacol. and Exp. Therap.
48(2): 241-265
- (555) Vojnovich, C., et al. 1960. A note on the enzymic
reduction of protein in high-amylose corn. Cereal
Chem. 37: 576-578
- (556) Waldschmidt-Leitz, E., and A. Harteneck. 1924. Zur
Kenntnis der spontanen Aktivierung des Trypsins.
IV. Zur Spezifität tierischer Proteasen. Ztschr.
f. physiol. Ch. 149: 221-30

- (557) Waldschmidt-Leitz, E., and G. Künstner. 1927.
Fraktionierte enzymatische Hydrolyse des Histons.
IV. über enzymatische Proteolyse. Ztschr. physiol.
Chem. 171: 290-315
- (558) Waldschmidt-Leitz, E., et al. 1933. Über den Einfluß
von Sulfhydrylverbindungen auf enzymatische Prozesse.
Hoppe-Seyler's Z. physiol. Chem. 214: 75-88
- (559) Wallenfels, K. 1950. Determination of proteolytic
activity of enzyme preparations. Critical comments
on the application of the Anson proteinase test.
Biochem. Z. 321: 189-196
- (560) Walton, R. P. 1931. Papain. Behavior in peritoneal
cavity. J. Pharmacol. and Exper. Therap. 43:
487-497
- (561) Walton, R. P., and C. M. Segural. 1933. Immunological
reaction to enzyme papain. Biochem. J. 26:
1750-1753
- (562) Wang, H., and N. Maynard. 1955. Enzymic tenderiza-
tion of meat. I. Basic technique and histological
observations of enzymic action. Food Research.
20: 587-597
- (563) Watson, A. F., et al. 1938. The enzyme papain and its
value in the preparation of some pharmaceutical
products. Quart. J. Pharm. and Pharmacol. 11(3):
391-400
- (564) Weir, C. E., et al. 1958. Enzymic tenderization of
meat. II. Panel and histological analyses of meat
treated with liquid tenderizers containing papain.
Food Research. 23: 411-422
- * (565) Wells, H., et al. 1965. Taste receptors and siala-
denotrophic action of proteolytic enzymes in rats.
Am. J. Physiol. 208(5): 877-881

- (566) Wenzel, J., and H. Arno. 1969. Enzyme-histochemical and morphological findings in the guinea pig myocardium following the application of various cardiotoxic substances. *Exp. Pathol.* 3(6): 305-320
- (567) Wenzel, J., and A. Hecht. 1969. Effects of cardiotoxic substances on the enzyme-histochemical picture of the heart in the guinea pig. *Beitr. Pathol. Anat. Allg. Pathol.* 138(3): 321-336
- (568) West German Patent. 1969. 1 442 333. Shaler, R. G., et al. Process for manufacturing fermentation liquids, in particular beer.
- (569) West German Patent. 1969. 1 442 337. Stone, I. M. Means for the cold stabilization of fermented malt drinks.
- (570) West German Patent. 1969. 1 492 682. Hogam, F. M. Enzyme-preparation for tenderizing meat.
- (571) West German Patent. 1969. 1 492 826. Kilpadi, V., et al. Process for the manufacturing of cheese.
- (572) West German Patent. 1970. Fieffe, C. F. Process for improving natural protein substances.
- (573) Westerborn, O. 1961. The effect of papain on epiphyseal cartilage. A morphological and biochemical study. *Acta Chir. Scand. Suppl.* 270: 1-84
- (574) Westerborn, O. 1965. The effects of papain on epiphyseal cartilage. *Struct. Function Connective Skeletal Tissue, Proc., St. Andrews, Scot.* 1964, 456-458
- (575) Whitaker, J. R. 1957. Assay and properties of commercial ficin. *Food Research.* 22: 468-478

- (576) Williamson, J. D. 1966. The agglutination of fowl red blood cells by the hemagglutinins of vaccinia and Cocksackie A7 viruses. J. Gen. Microbiol. 45(3): 451-458
- (577) Willcox, I. C. 1970. Beer Stabilization. Brewer's Guardian 99(10, Suppl.): 24-27
- (578) Willstätter, R., and W. Grassmann. 1928. Zur Freilegung des Invertins aus der Hefe. Biochem. Ztschr. 203: 308-312.
- (579) Witt, C. R., et al. 1970. Chillproofing beer with insoluble papain. Brewer's Digest. 45(10): 70, 72, 74, 76, 78, & 79
- (580) Wohlisch, E., and L. Juhling. 1938. Thrombin as a fibrinogen denaturase and its relation to papain. Biochem. Z. 297: 353-368
- (581) Wolf, E. P. 1922. Experimentelle Untersuchungen über Entzündung. II. Experimentelle chemische Entzündung in vivo. J. Exp. Med. 37: 511-24
- (582) Woodward, A. A. Jr. 1948. Protoplasmic clotting in isolated muscle fibers. J. Cell and Comp. Physiol. 31(3): 359-394
- * * (583) Wyzan, H. S., et al. 1951. The in vivo proteolytic activity of papain as demonstrated by nitrogen balance studies in dogs. Ann. N. Y. Acad. Sci. 54(2): 282-296
- (584) Yamashita, M., et al. 1970. Supplementing sulphur-containing amino acids by plastein reaction. Agricultural and Biological Chemistry. 34(10): 1593-1596
- (585) Yamashita, M., et al. 1971. Enzymatic modification of proteins in foodstuffs. V. Plastein yields and their pH dependence. Agricultural and Biological Chemistry 35(1): 86-91

- (588) Yaoi, H., and K. Nagase. 1928. Antigenic properties of papain digested bacterial antigens. *Scient. Rep. Gov. Inst. Infect. Dis.* 6: 127-137
- (589) Yoshinaka, R., and S. Ikeda. 1969. Effect of protease treatment on the physical properties of dough and gel of gluten. *Mem. Res. Inst. Food Sci. Kyoto Univ.* 30: 16-23
- (590) Yoshioka, M. 1951. The action of papain. *Bull. Inst. Chem. Research, Kyoto Univ.* 24: 79-80
- (591) Zacco, M., and G. Pratesi. 1947. Effects, immunologic reactions. *Boll. Soc. ital. biol. sper.* 23(4): 419-420
- (592) Zacco, M., and G. Pratesi. 1947. Histolytic effects of parenteral administration; experimental study. *Boll. Soc. ital. biol. sper.* 23: 548-550
- (593) Zacco, M., and G. Pratesi. 1947. Toxicity shock produced by papain. *Boll. Soc. ital. biol. sper.* 23: 551-552
- (594) Zacco, M., and G. Pratesi. 1947. Shock produced by parenteral administrations of papain corresponding with anaphylactic shock. *Boll. Soc. ital. biol. sper.* 23: 553-555
- (595) Zacco, M., and G. Biozzi. 1947. Effects. Anti-anaphylactic effects. 23: 792-793
- (596) Zacco, M. 1949. Histamine content of lungs after artificial pneumothorax. *Accad. Med.* 64: 387-388
- (597) Zakowski, J. 1931. Effect on urease preparations. *Ztschr. f. physiol. Chem.* 202: 249-267

- (598) Zakowski, J. 1933. Neue Versuche über die Einwirkung von Papain auf Ureasepräparate. Hoppe-Seyler's Z. physiol. Chem. 202: 249-267
- (599) Zalmanzon, E. S. 1949. Action of protease on nuclear inclusions in silkworm jaundice infections. Mikrobiologiya. 18: 361-365
- (600) Zappacosta, S., et al. 1968. Mechanism of cleavage of rabbit IgG in two stages by soluble papain and reducing agent. J. Immunol. 100(6): 1268-1276
- (601) Zwicker, H., et al. 1952. Studies on reactions of incomplete Rh antibodies with enzyme treated Rh positive cells. J. Immunol. 69(4): 415-421